



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2019

---

## **Multivariate meta-analyses of mitochondrial complex I and IV in major depressive disorder, bipolar disorder, schizophrenia, Alzheimer disease, and Parkinson disease**

Holper, L ; Ben-Shachar, D ; Mann, J John

**Abstract:** Complex I (NADH dehydrogenase, NDU) and complex IV (cytochrome-c-oxidase, COX) of the mitochondrial electron transport chain have been implicated in the pathophysiology of major psychiatric disorders, such as major depressive disorder (MDD), bipolar disorder (BD), and schizophrenia (SZ), as well as in neurodegenerative disorders, such as Alzheimer disease (AD) and Parkinson disease (PD). We conducted meta-analyses comparing complex I and IV in each disorder MDD, BD, SZ, AD, and PD, as well as in normal aging. The electronic databases Pubmed, EMBASE, CENTRAL, and Google Scholar, were searched for studies published between 1980 and 2018. Of 2049 screened studies, 125 articles were eligible for the meta-analyses. Complex I and IV were assessed in peripheral blood, muscle biopsy, or postmortem brain at the level of enzyme activity or subunits. Separate meta-analyses of mood disorder studies, MDD and BD, revealed moderate effect sizes for similar abnormality patterns in the expression of complex I with SZ in frontal cortex, cerebellum and striatum, whereas evidence for complex IV alterations was low. By contrast, the neurodegenerative disorders, AD and PD, showed strong effect sizes for shared deficits in complex I and IV, such as in peripheral blood, frontal cortex, cerebellum, and substantia nigra. Beyond the diseased state, there was an age-related robust decline in both complexes I and IV. In summary, the strongest support for a role for complex I and/or IV deficits, is in the pathophysiology of PD and AD, and evidence is less robust for MDD, BD, or SZ.

DOI: <https://doi.org/10.1038/s41386-018-0090-0>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-197233>

Journal Article

Accepted Version

Originally published at:

Holper, L; Ben-Shachar, D; Mann, J John (2019). Multivariate meta-analyses of mitochondrial complex I and IV in major depressive disorder, bipolar disorder, schizophrenia, Alzheimer disease, and Parkinson disease. *Neuropsychopharmacology*, 44(5):837-849.

DOI: <https://doi.org/10.1038/s41386-018-0090-0>

# Multivariate meta-analyses of mitochondrial complex I and IV in major depressive disorder, bipolar disorder, schizophrenia, Alzheimer disease and Parkinson disease

Running title: Meta-analyses of mitochondrial complex I and IV

Holper L <sup>1</sup>, Ben-Shachar D <sup>2</sup>, Mann JJ <sup>1</sup>

<sup>1</sup> Division of Molecular Imaging and Neuropathology, Columbia University and New York State Psychiatric Institute, New York, NY, USA

<sup>2</sup> Laboratory of Psychobiology, Department of Psychiatry, Rambam Health Care Campus, Rappaport Faculty of Medicine, Technion IIT, Haifa, Israel

Corresponding author: Lisa Holper, Division of Molecular Imaging and Neuropathology, Columbia University and New York State Psychiatric Institute, 1051 Riverside Drive, New York, NY 10032, US, Phone: 646-774-7637, Mail: [holper@ini.phys.ethz.ch](mailto:holper@ini.phys.ethz.ch)

## **Abstract**

Complex I (NADH dehydrogenase, NDU) and complex IV (cytochrome-c-oxidase, COX) of the mitochondrial electron transport chain have been implicated in the pathophysiology of major psychiatric disorders such as major depressive disorder (MDD), bipolar disorder (BD) and schizophrenia (SZ), as well as in neurodegenerative disorders, such as Alzheimer disease (AD) and Parkinson disease (PD).

We conducted meta-analyses comparing complex I and IV in each disorder MDD, BD, SZ, AD PD, as well as in normal aging. The electronic databases Pubmed, EMBASE, CENTRAL and Google Scholar, were searched for studies published between 1980 and 2018. Of 2049 screened studies, 125 articles were eligible for the meta-analyses. Complex I and IV were assessed in peripheral blood, muscle biopsy, or postmortem brain at the level of enzyme activity or subunits.

Separate meta-analyses of mood disorder studies, MDD and BD, revealed moderate effect sizes for similar abnormality patterns in the expression of complex I with SZ in frontal cortex, cerebellum and striatum, whereas evidence for complex IV alterations was low. By contrast, the neurodegenerative disorders, AD and PD, showed strong effect sizes for shared deficits in complex I and IV such as in peripheral blood, frontal cortex, cerebellum and substantia nigra. Beyond the diseased state, there was an age-related robust decline in both complexes I and IV. In summary, the strongest support for a role for complex I and/or IV deficits, is in the pathophysiology of PD and AD, and evidence is less robust for MDD, BD or SZ.

## Introduction

Mitochondrial dysfunction is implicated in the pathophysiology of major psychiatric disorders, such as major depressive disorder (MDD) (Bansal and Kuhad, 2016), bipolar disorder (BD) (Kato, 2017) and schizophrenia (SZ) (Bergman and Ben-Shachar, 2016), as well as neurodegenerative disorders, such as Alzheimer disease (AD) (Onyango *et al*, 2017) and Parkinson disease (PD) (Onyango *et al*, 2017). Mitochondria are intracellular organelles that produce adenosine triphosphate (ATP), the main source of cellular energy. Impaired mitochondrial function results in decreased ATP production, impaired bioenergetics, apoptosis and oxidative stress (Hroudová and Fišar, 2011). Prior to the generation of ATP, mitochondria direct electrons extracted from nutrients into a transmembrane proton gradient and this process is mediated by the electron transport chain (ETC).

Research has identified two enzymes of the ETC located at the inner mitochondrial membrane as being particular impaired in these five disorders MDD, BD, SZ, AD and PD. The first enzyme, complex I (NADH dehydrogenase, NDU) consists of 45 subunits, seven of which are encoded by mitochondrial DNA (mtDNA) and the remaining subunits by nuclear DNA (nDNA). Complex I is one of the entry enzymes of cellular respiration or oxidative phosphorylation in the mitochondrion. It is also the largest multimeric enzyme complex of the ETC and is a major contributor to the generation of the proton gradient across the mitochondrial inner membrane, which drives ATP production. The second enzyme of interest, complex IV (cytochrome-c-oxidase, COX) consists of 13 subunits, three of which are encoded by mtDNA, the remainder by nDNA. Complex IV catalyzes the final step in the mitochondrial ETC and, due to its rate-limiting role in this oxidative process (Arnold, 2012), has been proposed as a key markers of mitochondrial function (Srinivasan and Avadhani, 2012). Numerous excellent reviews (Bansal and Kuhad, 2016; Bergman and Ben-Shachar, 2016; Kato, 2017; Onyango *et al*, 2017) have discussed the details of impairments in both

complex I and IV enzyme activities and subunit assembly within each of the above mentioned disorders. A meta-analysis summarizing the findings across these disorders could not be found in the literature. The remaining complexes II (succinate dehydrogenase ), III (cytochrome c reductase) and V (ATP synthase) either have not been studied, or they have been studied to a much smaller degree in these five disorders, compared with complex I and IV, and thus there are too few data for a meta-analysis.

We chose these five disorders MDD, BD, SZ, AD and PD not only because of the potential common mitochondrial dysfunction, but also based on their clinical similarities. Though regarded as different disorders in major classification systems like DSM and ICD, there is also overlap in clinical symptoms. Depression is found in mood disorders, but is also frequent in schizophrenia and both AD and PD (Lancôt *et al*, 2017; Zhuo *et al*, 2017). Psychotic symptoms are observed in MDD and BD as well as SZ ((DSM-IV-TR), 2000). Although Alzheimer's and Parkinson's diseases have distinct brain histopathology, both are age-related neurodegenerative conditions characterized by memory loss and depression and have some commonality in molecular pathogenesis, such as proteinopathy, neuroinflammation and oxidative stress (Aarsland *et al*, 2001). Cross-disorder commonalities are also found in schizophrenia sharing dopaminergic abnormalities with Parkinson's (Brisch *et al*, 2014) and cognitive impairment (dementia praecox) with Alzheimer's (Lyketsos CG and Peters ME, 2015).

Beyond disease states, normal aging plays an independent role in mitochondrial ETC function (Sun *et al*, 2016). To separate the influence of age from that of disease, we therefore additionally meta-analyzed complex I and IV functioning in healthy older versus younger individuals. The present meta-analyses aimed to provide a synthesis of the work done on complex I and IV that can guide future research and stimulate development of novel *in-vivo* technologies such as for example the assessment of the redox states of complex I

NAD<sup>+</sup>/NADH ratio (Blacker and Duchen, 2016; Zhu *et al*, 2015) or oxidized complex IV (Bale *et al*, 2016).

## **Materials and methods**

### **Literature search and study identification**

We conducted a structured literature search in PubMed, EMBASE, CENTRAL and Google Scholar, to identify studies published between January 1980 and January 2018 using the search strings ‘NADH dehydrogenase’ OR ‘cytochrome-c-oxidase’ OR ‘complex I’ OR ‘complex IV’ AND ‘major depressive disorder’ OR ‘bipolar disorder’ OR ‘schizophrenia’ OR ‘Alzheimer’ OR ‘Parkinson’ OR ‘age/aging’. We manually reviewed reference lists in all retrieved articles for related publications.

Inclusion criteria were: studies investigated either complex I and/or complex IV, studies published in the English language, studies reporting original human data and studies that investigated patients in comparison with a control group. Exclusion criteria were: animal studies, cellular studies, genetic studies, studies with less than three subjects per group, case reports, letters to the editor and editorials and publications not reporting original data.

### **Data extraction**

Aggregated data were extracted for each of the disorders MDD, BD, SZ, AD, PD and normal aging (AGING). Aggregated data contained quantitative data (i.e., number of patients and controls, mean age of patients and controls), qualitative information (i.e., methods such as enzyme activity or subunits at the level of mRNA or protein expression, and tissues assayed

such as peripheral blood, muscle or brain regions of interest, ROIs) and outcomes (effect sizes in terms of the standardized mean difference, SMD, and p-values). Enzyme activity was extracted as a corrected ratio to citrate synthase to normalize for mitochondrial content, if available. For studies on AGING, we chose a breakpoint of mean 60 years (Cabr   *et al*, 2017) to derive group differences between older (> 60) and younger (< 60) individuals using t-test. In case of medication reported in the studies, both un-medicated and medicated cohorts were included. In case of BD, data were included from both BD I and BD II during depressive episodes, but not manic episodes.

The authors of some studies were contacted for permission to reuse original data (Altar *et al*, 2005; Karry *et al*, 2004; Konradi C *et al*, 2004; Manczak *et al*, 2004; Thomas *et al*, 2012; Washizuka *et al*, 2005, 2009); no answer was received in some cases and the data could therefore not or only partially included (Altar *et al*, 2005; Konradi C *et al*, 2004; Manczak *et al*, 2004; Thomas *et al*, 2012). Some data were read from figures (Andreazza *et al*, 2013; Andreazza AC *et al*, 2010; Cabr   *et al*, 2017; Chagnon *et al*, 1995; Chandrasekaran *et al*, 1998; Devi *et al*, 2006; Distefano *et al*, 2017; Gatt *et al*, 2016; Gueugneau *et al*, 2015; Hsieh *et al*, 1994; Kim *et al*, 2016; Kish *et al*, 1992; Mehler-Wex *et al*, 2006; Munkholm *et al*, 2015; Mythri *et al*, 2011; Ojaimi *et al*, 1999; Parker *et al*, 2008; Safdar *et al*, 2010; Simonian and Hyman, 1993, 1995; Trounce *et al*, 1989).

## **Multivariate random-effect meta-analyses**

Separate meta-analyses were first computed *for each disorder*. Moderator variables were generated representing the combination of two aspects: 1) method (enzyme activity or subunit); and 2) tissue (blood, muscle or ROI). A moderator was included in the meta-analysis only if it was reported by at least three studies. This threshold was chosen after extensive prior testing in order to avoid spurious results.

A summary meta-analysis was computed *across all disorders*. Here, the moderator variable consisted of the five disorders MDD, BD, SZ, AD, PD and AGING. This analysis included all data found to be eligible in the present work and thus also included rare outcomes that could not be included in the separate disease/aging meta-analyses.

For all meta-analyses, a multivariate random-effect model was applied based on the Metafor package (Viechtbauer, 2010) as implemented in R (R Development Core Team, 2008). The multivariate model accounts for heterogeneity and dependency in the underlying true effects of multiple moderators that can overlap within subjects and studies (random factors). To adjust for age effects, the age-difference between patients and controls was added as a continuous covariate. To adjust for sample size, the effects were weighted based on study size. To allow for heterogeneity differences between moderators, an unstructured variance-covariance matrix was applied (`rma.mv` function; observed effects = SMD + age-difference; variance-covariance matrix = COV; weight = study size; variance structure = UN; mods = moderators/disorder; random factors = subject/study; method = REML).

Forest plots were generated to illustrate the results of the multivariate models in terms of the weighted beta coefficients with 95% confidence intervals (CIs) and p-values. Heterogeneity was assessed using Cochran's Q-test and the inconsistency  $I^2$  statistic that directly indicates to what extent each outcome contributes to the total variance. Publication bias was assessed using Egger's regression analysis.



## Results

### Data extraction

Of 2049 screened studies, 125 studies were eligible for the meta-analyses, some of them for more than one disorder (MDD N = 7, BD N = 19, SZ N = 21, AD N = 43, PD N = 35, AGING N = 18) (**Table 1**). The overall number of studies reporting on complex I (total N = 101, MDD N = 6, BD N = 19, SZ N = 15, AD N = 19, PD N = 32, AGING N = 11) and complex IV (total N = 104, MDD N = 4, BD N = 10, SZ N = 10, AD N = 39, PD N = 28, AGING N = 15) were similar but varied between disorders. Both complex I and IV were analysed in either peripheral blood elements, such as platelets, lymphocytes or other blood cells (MDD N = 1, BD N = 10, SZ N = 8, AD N = 9, PD N = 14, AGING N = 2), in muscle biopsies (MDD N = 2, BD N = 0, SZ N = 0, AD N = 0, PD N = 9, AGING N = 14), or in postmortem brain samples (MDD N = 4, BD N = 8, SZ N = 13, AD N = 34, PD N = 14, AGING N = 2). Both complex I and IV were analysed at the level of enzymatic activity (either in blood, muscle, or brain, MDD N = 4, BD N = 5, SZ N = 11, AD N = 32, PD N = 32, AGING N = 14), or of their subunits by mRNA level (only brain, MDD N = 3, BD N = 7, SZ N = 7, AD N = 7, PD N = 1, AGING N = 2) or protein level (only brain, MDD N = 3, BD N = 12, SZ N = 7, AD N = 6, PD N = 3, AGING N = 5). In the following, enzyme activity is denoted with NDU/COX respectively, whereas subunits as assessed by either mRNA or protein expression are referred to with their full names.

Note that we combined data from levels of mRNA/protein expression for all subunits. The correlation between mRNA and protein expression is generally thought to be around 40% (Abreu et al, 2009; Vogel and Marcotte, 2012), which is typically attributed to other levels of regulation between transcript and protein product (**Maier et al, 2009**). We separated the

findings in a pre-analysis but found no significant differences between mRNA and protein expression and therefore presented the results together.

Assay of enzyme activity was more common in studies of neurodegenerative disorders, AD (88%) and PD (66%), whereas the assessment of mRNA/protein expression was more common in studies of mood disorders, MDD (60%), BD (79%) and SZ (56%). None of the five disorders, however, showed differences in enzyme activity or mRNA/protein expression for complex I or IV ( $p > 0.05$ ). Moreover, there were no differences between peripheral blood and brain in the enzyme activity/mRNA/protein levels of complex I and IV ( $p > 0.05$ ).

All studies involved subjects over 18 years old, except two studies examined early-onset SZ patients below age 18 years (Mehler-Wex *et al*, 2006; Taurines *et al*, 2010). The mean age-difference between cases and controls differed between disorders (main effect  $F = 35.21$ ,  $p < 0.0001$ ). Psychiatric studies, MDD (mean age-difference  $\pm$  STD =  $-0.7 \pm 3.9$ ), BD (mean age-difference  $\pm$  STD =  $0.8 \pm 4.9$ ) and SZ (mean age-difference  $\pm$  STD =  $-1.0 \pm 3.7$ ) included closer age-matched cases and controls compared with AD (mean age-difference  $\pm$  STD =  $5.0 \pm 7.5$ ) or PD (mean age-difference  $\pm$  STD =  $5.3 \pm 5.9$ ) as indicated by significant post-hoc comparisons ( $p < 0.0001$ ). Some studies did not provide mean ages for cases and/or controls (Cooper *et al*, 1993; DiDonato *et al*, 1993; Hirai *et al*, 2001; Reichmann *et al*, 1993; Wiedemann *et al*, 1999), for which we therefore assumed zero age-difference.

## **Multivariate random-effect meta-analyses**

Meta-analyses were performed for each of the disorders MDD, BD, SZ, AD, PD and AGING. Based on the precondition that a moderator was reported at least three times across studies, the number of studies finally included in the meta-analyses (and the percentage of the eligible studies) differed between disorders MDD (complex I  $N = 4$  (67%), complex IV  $N = 2$  (50%)),

BD (complex I N = 16 (84%), complex IV N = 3 (30%)), SZ (complex I N = 14 (93%), complex IV N = 6 (60%)), AD (complex I N = 12 (63%), complex IV N = 33 (85%)), PD (complex I N = 29 (91%), complex IV N = 24 (86%) and AGING (complex I N = 7 (64%), complex IV N = 14 (93%)).

## Mood disorders

In MDD (**Figure 1a**), complex I subunits NDUFS1<sub>cerebellum</sub> ( $p = 0.0075$ ), NDUFV1<sub>cerebellum</sub> ( $p = 0.00014$ ) and NDUFV2<sub>cerebellum</sub> ( $p = 0.00051$ ) were lower in cerebellum and subunit NDUFV1<sub>frontal</sub> ( $p = 0.0047$ ) was lower in frontal cortex compared with controls. No differences were found for complex IV (note that only in this case, we set the moderator threshold to at least two).

In BD (**Figure 1b**), complex I subunit NDUFS1<sub>cerebellum</sub> ( $p < 0.0001$ ) was lower in cerebellum, subunits NDUFS1<sub>frontal</sub> ( $p = 0.033$ ) and NDUFV7<sub>frontal</sub> ( $p < 0.0001$ ) were lower in frontal cortex and subunit NDUFS1<sub>striatum</sub> ( $p = 0.0023$ ) was lower in striatum compared with controls. No differences were found for complex IV.

In SZ (**Figure 1c**), complex I subunit NDUFV2<sub>blood</sub> ( $p = 0.019$ ) was lower in blood, whereas subunit NDUFS1<sub>blood</sub> ( $p = 0.00086$ ) was higher in blood compared with controls. Further, subunits NDUFS1<sub>striatum</sub> ( $p = 0.00072$ ), NDUFV1<sub>striatum</sub> ( $p < 0.0001$ ) and NDUFV2<sub>striatum</sub> ( $p < 0.0001$ ) were lower in striatum, subunit NDUFV1<sub>frontal</sub> ( $p = 0.011$ ) was lower in frontal cortex, whereas subunit NDUFV2<sub>parietal</sub> ( $p = 0.019$ ) was higher in parietal cortex compared with controls. Complex IV enzyme activity was lower in frontal cortex (COX<sub>frontal</sub>  $p = 0.016$ ), but higher in basal ganglia (COX<sub>NA/GP/P</sub>  $p < 0.0001$ , nucleus accumbens, globus pallidus and putamen) compared with controls.

We also assessed the overlap in findings between MDD, BD and SZ using meta-regressions. Results confirmed lower NDUFS1/NDUFV1/NDUFV2 levels in the cerebellum in MDD ( $p =$

0.005/ $p = 0.001/p = 0.006$ ) and in some subunits also BD ( $p = 0.088/p = 0.001/p = 0.258$ ) compared to SZ, respectively. By contrast, lower NDUFS1/NDUFV1/NDUFV2 levels in striatum were observed in SZ compared to MDD ( $p = 0.001/p < 0.0001/p < 0.0001$ ) and in some subunits also BD ( $p = 0.250/p < 0.0001/p < 0.0001$ ), respectively. Further, both BD ( $p = 0.024$ ) and SZ ( $p = 0.008$ ) revealed higher levels of NDUFV2 in the parietal cortex compared to MDD. Last, in BD, subunit NDUFS1<sub>blood</sub> levels ( $p = 0.002$ ) in blood and subunit NDUFS7<sub>frontal</sub> ( $p = 0.009$ ) in frontal cortex were lower compared to SZ.

### Neurodegenerative disorders

In AD (**Figure 2a**), complex I enzyme activity in the temporal cortex (NDU<sub>temporal/entorhinal</sub>  $p = 0.0023$ ) and subunits NDUF1/4/7-9<sub>blood</sub> ( $p < 0.0001$ ), NDUF2/3/6<sub>blood</sub> ( $p < 0.0001$ ) and NDUF3/4/5<sub>blood</sub> ( $p < 0.0001$ ) were lower in blood compared with controls. It should be noted that these subunits were reported by Lunnon et al. (Lunnon *et al*, 2012), which was the largest study eligible for the present work with a total of 209 subjects (patients  $N = 104$ , controls  $N = 105$ ). Initially only a small subset of that study could be included in the meta-analysis because of the high moderator threshold; in order to include more of the results from that study, we summarized results across subunits to build sufficient moderators.

Complex IV enzyme activity was lower in frontal cortex (COX<sub>frontal</sub>  $p = 0.044$ ), motor cortex (COX<sub>motor</sub>  $p = 0.0003$ ), occipital cortex (COX<sub>occipital</sub>  $p = 0.032$ ), parietal cortex (COX<sub>parietal</sub>  $p = 0.00069$ ), temporal/entorhinal cortex (COX<sub>temporal/entorhinal</sub>  $p = 0.00012$ ) and hippocampus (COX<sub>hippocampus</sub>  $p < 0.0001$ ) compared to controls. Subunit COX2<sub>hippocampus</sub> in hippocampus ( $p < 0.0001$ ) and subunit COX7<sub>blood</sub> in peripheral blood ( $p = 0.011$ ) were lower.

In PD (**Figure 2b**), complex I enzyme activity in peripheral muscle (NDU<sub>muscle</sub>  $p = 0.000084$ ) as well as in substantia nigra (NDU<sub>substantia nigra</sub>  $p = 0.006$ ) was lower compared to controls. Complex IV enzyme activity in frontal cortex (COX<sub>frontal</sub>  $p = 0.036$ ) was also lower.

The overlap between AD and PD was assessed using meta-regressions. Results showed comparable patterns of enzyme activity in blood ( $\text{COX}_{\text{blood}}$   $p = 0.981$ ) and brain such as in frontal cortex ( $\text{NDU}_{\text{frontal}}$   $p = 0.171$ ,  $\text{COX}_{\text{frontal}}$   $p = 0.711$ ), cerebellum ( $\text{COX}_{\text{cerebellum}}$   $p = 0.582$ ) and substantia nigra ( $\text{COX}_{\text{substantia nigra}}$   $p = 0.671$ ), indicating no significant differences between AD and PD.

## **Aging**

To assess complex I and IV functioning in normal aging, we performed a separate meta-analysis in healthy subjects using a breakpoint of 60 years (Cabr  *et al*, 2017) (**Figure 2c**). Older ( $\geq 60$  years) subjects in comparison with younger ( $< 60$  years) subjects had lower complex I enzyme activity in frontal cortex ( $\text{NDU}_{\text{frontal}}$   $p < 0.0001$ ) and muscle ( $\text{NDU}_{\text{muscle}}$   $p < 0.0001$ ,  $\text{NDUFB8}_{\text{muscle}}$   $p = 0.00015$ ). Similarly, complex IV enzyme activity was lower in frontal cortex ( $\text{COX}_{\text{frontal}}$   $p = 0.00072$ ) and muscle ( $\text{COX}_{\text{muscle}}$   $p = 0.00027$ ,  $\text{COX2}_{\text{muscle}}$   $p < 0.0001$ ) in older compared to younger subjects.

## **Summary comparison of all cases and controls**

A summary meta-analysis was performed to categorize the disorders according to the severity of overall mitochondrial impairments (**Figure 3**). Robust deficits were observed for complex I in PD ( $p = 0.0016$ ) and AGING ( $p = 0.009$ ) and for complex IV in AD ( $p = 0.004$ ) and AGING ( $p < 0.0001$ ). MDD ( $p < 0.0001$ ) also showed an effect on complex I; however, it was based on small samples sizes from few studies and can therefore not be considered a robust result. The main effect across both complex I and IV was significant in PD ( $p = 0.00058$ ), AD ( $p = 0.049$ ) and AGING ( $p < 0.0001$ ) indicating that these three conditions were also affected if both complex I and IV are considered.

## Heterogeneity and publication bias

Overall, there was a high degree of heterogeneity (**Table 2**). The  $I^2$  for the five disorders MDD, BD, SZ, AD and PD ranged between 80-100%. Noticeable, there was low heterogeneity in AGING in both complex I ( $Q = 27.934$ ,  $p = 0.218$ ,  $I^2 = 18\%$ ) and IV ( $Q = 11.684$ ,  $p = 1$ ,  $I^2 = 0\%$ ). Putative low heterogeneity was observed in MDD for complex IV ( $Q = 0.030$ ,  $p = 1$ ,  $I^2 = 0\%$ ), most likely explained by the fact that the data came from few studies ( $N = 4$ ).

Publication bias as assessed using Egger's regression test (**Table 2**) provided non-significant results for most of the data indicating no publication bias. Considerable publication bias was only found in PD for complex I ( $p = 0.043$ ) and in AGING for complex IV ( $p = 0.014$ ).

## Discussion

Our findings provide a picture of the current findings on complex I and IV in mood and neurodegenerative disorders (**Figure 4**). We found that mood disorders, MDD and BD, share commonalities with SZ in having low expression of complex I subunits. Neurodegenerative disorders, AD and PD, share similarities in complex I and IV enzyme activity abnormalities. Due to heterogeneity in study findings, strong evidence for complex I and/or IV deficits is present for AD and PD, whereas the overall proof for the involvement of these complexes in MDD, BD and SZ is less robust and requires further research. Normal aging is a substantial independent factor contributing to both complex I and IV decline and should be considered when interpreting study heterogeneity.

### Mood disorders

Mood disorders show shared patterns of complex I subunits NDUFS1, NDUFV1 and NDUFV2 between MDD and BD on the one hand, and between BD and SZ on the other hand (**Figure 1**) (Andreazza *et al*, 2013; Andreazza AC *et al*, 2010; Ben-Shachar and Karry, 2008; Iwamoto *et al*, 2005; Karry *et al*, 2004; Kim *et al*, 2016; Sun *et al*, 2006). While, MDD and to some degree BD, have lower levels of NDUFS1, NDUFV1, NDUFV2 in cerebellum compared with SZ, the latter had lower NDUFS1, NDUFV1, NDUFV2 levels in striatum compared with MDD and to some degree BD; this indicates a shared mechanism between MDD and BD that is distinct from SZ. By contrast, BD and SZ had elevated levels of NDUFV2 in parietal cortex compared with MDD, which may indicate a shared mechanism between BD and SZ. These brain regions, i.e., the frontal cortex (Brakowski *et al*, 2017; Hibar *et al*, 2017), cerebellum (Jiang *et al*, 2017; Shinn *et al*, 2017) and striatum (Pacifico and Davis, 2016), have been previously suggested to contribute to the pathophysiology of mood disorders and schizophrenia spectrum (Dietsche *et al*, 2017). Distinct from the mood

disorders, SZ has higher complex IV levels in basal ganglia (Prince *et al*, 1999, 2000) perhaps related to previous neuropathological findings in putamen, nucleus accumbens and globus pallidus (Mamah *et al*, 2007; Womer *et al*, 2014).

Notably, while most of the eligible studies pointed to dysregulation of complex I and/or IV, the reported direction of change has not always been consistent. While some of the heterogeneity may be due to potential differences in the clinical picture like presence of psychosis in MDD or BD and presence of depression in SZ, several other explanations may underlie these discrepancies. First, there may be state-related differences in the expression of complex I and IV. For example BD patients showed up-regulation of complex I subunits during manic compared to depressive episodes (Akarsu *et al*, 2015) maybe related to elevated brain metabolism in frontal and temporal cortex (Shao *et al*, 2008). Similarly in SZ, state-dependence has been suggested with increased complex I activity in active psychotic patients but decreased activity in patients with residual schizophrenia (Ben-Shachar *et al*, 2007; Dror *et al*, 2002; Rosenfeld *et al*, 2011). Unfortunately, postmortem studies were unable to provide information on whether patients were depressed, manic or euthymic at time of death. Second, there may be brain region-specific heterogeneity, possibly indicating that ETC energy production varies from region to region, as previously exemplarily shown (Ben-Shachar and Karry, 2008). Third, there may be tissue-specific differences between blood samples from living patients and brain samples obtained postmortem. For example, it has been hypothesized that ETC genes are up-regulated in blood, but down-regulated in some or all brain regions in BD (Beech *et al*, 2010) and SZ (Ben-Shachar and Karry, 2007), although this was not confirmed statistically by the present meta-analysis. Fourth, the majority of patients were receiving psychotropic medication, such as antidepressants, mood stabilizers and antipsychotics. All of these pharmacological agents have been reported to potentially interfere or even inhibit the mitochondrial ETC (Manji *et al*, 2012). Finally, differences in brain pH at



the time of death may explain difference in gene expression (Vawter *et al*, 2006). Larger sample studies of medication-free patients are needed to clarify more precisely the relationships between diagnosis, mood state, treatment status and brain pH in mood disorders and schizophrenia pathophysiology. [There is also evidence of shared global gene expression patterns between mood disorders and schizophrenia that go beyond purely mitochondrial function that could be investigated in relation \(Gandal \*et al\*, 2018\).](#)

## Neurodegenerative disorders

Neurodegenerative disorders AD and PD show similarities in the impairment of complex I and IV enzyme activities in the blood, frontal cortex, cerebellum and substantia nigra (**Figure 2**). These brain regions are potentially involved in the cognitive decline and neuronal loss in AD (Boublay *et al*, 2016; Perluigi *et al*, 2016) as well as in the motor and non-motor related degeneration of neurotransmitter systems in PD (Atkinson-Clement *et al*, 2017; Gao and Wu, 2016).

In AD, there are a well-documented deficit in both complex I and IV (Coskun *et al*, 2012; Giachin *et al*, 2016; Onyango *et al*, 2017). These deficits are thought to be due to the neuronal toxicity induced by amyloid  $\beta$  peptide ( $A\beta$ ), an important component of AD pathogenesis, in that  $A\beta$  accumulates in mitochondria, directly inhibits mitochondrial enzymes, perpetuates oxidative stress, and leads to a hypometabolic state, which causes mitochondrial dysfunction (Cenini *et al*, 2016; Picone *et al*, 2014; Pinho *et al*, 2014; Readnower *et al*, 2011). The meta-analysis supported these findings in terms of down-regulated complex I and IV in the blood (Cardoso *et al*, 2004; Feldhaus *et al*, 2011; Fišar *et al*, 2016; Lunnon *et al*, 2012; Mancuso *et al*, 2003; Parker *et al*, 1990, 1994a; Sheehan *et al*, 1997; Valla *et al*, 2006), frontal cortex (Cavelier *et al*, 1995; Devi *et al*, 2006; Hirai *et al*, 2001; Kish *et al*, 1992; Long *et al*, 2012; Maurer *et al*, 2000; Parker *et al*, 1994b; Pérez-Gracia *et al*, 2008; Wong-Riley *et al*, 1997),

motor cortex (Bosetti *et al*, 2002; Valla *et al*, 2001; Wong-Riley *et al*, 1997), occipital cortex (Kish *et al*, 1992, 1999; Mutisya *et al*, 1994), parietal cortex (Chagnon *et al*, 1995; Kish *et al*, 1992, 1999; Mutisya *et al*, 1994; Reichmann *et al*, 1993; Wong-Riley *et al*, 1997), temporal/entorhinal cortex (Alikhani *et al*, 2011; Chandrasekaran *et al*, 1998; Cooper *et al*, 1993; Gu *et al*, 1998; Kish *et al*, 1992, 1999; Maurer *et al*, 2000; Mutisya *et al*, 1994; Reichmann *et al*, 1993; Wong-Riley *et al*, 1997) and hippocampus (Aksenov *et al*, 1999; Bosetti *et al*, 2002; Chagnon *et al*, 1995; Chandrasekaran *et al*, 1994; Cottrell *et al*, 2001; Kish *et al*, 1992; Maurer *et al*, 2000; Reichmann *et al*, 1993; Simonian and Hyman, 1993, 1995; Verwer *et al*, 2000; Wong-Riley *et al*, 1997). The strength of the effect sizes are in line with findings of neuronal loss in entorhinal cortex, hippocampus and association neocortex (Perluigi *et al*, 2016). Due to limited data, the meta-analysis was not able to distinguish between Braak stages in AD. A large study in AD (N = 148) reported less expression of complex I and IV subunits in entorhinal and frontal cortex in disease stages V-VI but not in stages I-II (Armand-Ugon *et al*, 2017). A smaller study (N = 18) compared complex I and IV enzyme activity in frontal cortex in stages III-VI to stages I-II but found no stage-dependence (Manczak *et al*, 2004). There were also no differences between early- versus late-onset AD (age < 60 versus age > 60) (Cardoso *et al*, 2004).

In PD, the meta-analysis points to down-regulation of both complex I and IV enzyme activity in the blood, muscle and brain. For example, inhibition of complex I has been reported in platelets (Benecke *et al*, 1993; Blake *et al*, 1997; Bravi *et al*, 1992; Gu *et al*, 1998; Krige *et al*, 1992; Parker *et al*, 1989; Varghese *et al*, 2009; Yoshino *et al*, 1992) and skeletal muscles (Bindoff *et al*, 1991; Blin *et al*, 1994; Cardellach *et al*, 1993; Nakagawa-Hattori *et al*, 1992; Shoffner *et al*, 1991), although there were dissenting studies in platelets (Bronstein *et al*, 2015; Martín *et al*, 1996) or muscle (Anderson *et al*, 1993; DiDonato *et al*, 1993; Mann *et al*, 1992). Similarly, complex IV has been found to be decreased in blood (Benecke *et al*, 1993; Bravi *et*

*al*, 1992; Haas *et al*, 1995; Krige *et al*, 1992; Parker *et al*, 1989; Shinde and Pasupathy, 2006) and muscle (Anderson *et al*, 1993; Bindoff *et al*, 1991; Cardellach *et al*, 1993; Shoffner *et al*, 1991), while others could not confirm this for either platelets (Blake *et al*, 1997; Hanagasi *et al*, 2005) or muscle (Blin *et al*, 1994; DiDonato *et al*, 1993; Mann *et al*, 1992; Nakagawa-Hattori *et al*, 1992; Wiedemann *et al*, 1999). Within the brain, some studies reported lower complex I enzyme activity in substantia nigra (Gu *et al*, 1998; Mann *et al*, 1992; Mizuno *et al*, 1990; Schägger, 1995; Schapira *et al*, 1990b) as well as low activity and impaired assembly of complex I in frontal cortex (Barroso *et al*, 1993; Gatt *et al*, 2016; Keeney *et al*, 2006; Mizuno *et al*, 1990; Mythri *et al*, 2011; Parker *et al*, 2008), the latter not being confirmed in the meta-analysis. Similarly, complex IV in the brain is reported to be low in frontal cortex (Barroso *et al*, 1993; Chagnon *et al*, 1995; Keeney *et al*, 2006; Mizuno *et al*, 1990; Parker *et al*, 2008) and substantia nigra (Chagnon *et al*, 1995; Gu *et al*, 1998; Mann *et al*, 1992; Schägger, 1995; Schapira *et al*, 1990a), the latter finding not confirmed here. Note that although all data eligible for the meta-analysis were obtained from patients suffering from the more common sporadic form of PD, derangements in mitochondrial metabolism and oxidative stress have also been strongly linked to the less common familial form of PD (Ryan *et al*, 2015).

Likewise for mood disorders, medication may affect mitochondrial functioning in neurodegeneration. Cholinesterase (ChE) inhibitors and amyloid-beta binding alcohol dehydrogenase (ABAD) modulators are used to treat the key symptoms in AD in order to improve activities of daily living, behavior and cognition (Grossberg, 2003). Most of these drugs can negatively affect both complex I and IV (Hroudová *et al*, 2017). There are however newly developed ChEs and ABADs as well as other more recent drug developments that target multiple AD pathophysiological pathways that are thought to have a better outcome on mitochondrial function (Korábečný *et al*, 2018). By contrast, in PD, levodopa, cardidopa, and

selegiline, which are used to treat PD motor symptoms, have not been shown not affect complex I and IV in PD (Dixit et al, 2013; Shults et al, 1995). Similarly, melatonin that is used as sleep/wake regulator in PD, acts as an effective antioxidant and mitochondrial function protector (Srinivasan et al, 2011). These observations support the hypothesis that impaired complex I and IV activity in PD patients is a characteristic of the disease and not due to medications.

## Aging

Normal aging has been associated with a decline in mitochondrial quality (Sun *et al*, 2016) and activity of both complex I and IV (Arnold, 2012; Navarro and Boveris, 2007). The meta-analysis supported these findings in terms of decreased levels of enzyme activity in the muscle (Boffoli *et al*, 1996; Distefano *et al*, 2017; Emelyanova *et al*, 2017; Gueugneau *et al*, 2015; Hsieh *et al*, 1994; Lanza *et al*, 2008; Ogborn *et al*, 2015; Pestronk *et al*, 2017; Rooyackers *et al*, 1996; Safdar *et al*, 2010; Zucchini *et al*, 1995) and frontal cortex (Boffoli *et al*, 1994, 1996; Cabré *et al*, 2017; Ojaimi *et al*, 1999) (**Figure 2c**). Across all meta-analyses, the effect of aging was the most robust both in terms of the strength of the summary effect sizes (**Figure 3**) as well as the low degree of heterogeneity (**Table 2**).

The decline in both complex I and IV in normal aging (Arnold, 2012; Navarro and Boveris, 2007) as indicated by this meta-analysis can represent a functionally serious impairment of mitochondrial function with respect to basal ATP production (Boveris *et al*, 1999). Therefore, aged neurons with lower mitochondrial mass and enzyme activities as well as more dysfunctional mitochondria may be unable to respond adequately to any increased ATP demand (Navarro and Boveris, 2004).

The term normal aging should be considered carefully. Normal aging is thought to be a function of both chronological age as well as biological age (Safdar et al, 2010). Although the

underlying mechanism is still controversial, age-dependent mitochondrial decline may contribute to loss of physical capacity and mental exertion (Christian and Shadel, 2014; Payne and Chinnery, 2015). The ‘free radical’ theories of aging (Harman, 1992; Miquel *et al*, 1980) state that reactive oxygen species (ROS) produced aberrantly during mitochondrial electron transport damage mitochondrial components, including mtDNA. Since, mtDNA encodes essential complex I and IV proteins (Bonawitz *et al*, 2006), its damage and mutagenesis would disrupt ETC complex assembly and, in turn, lead to more ROS production, cellular oxidative stress and tissue dysfunction that promote aging (Mandavilli *et al*, 2002). This theory, however, has been thought to be too simplistic and there is debate about other age-dependent cellular changes that may be as important in precipitating the decline in mitochondrial ETC (Christian and Shadel, 2014).

### **Summary comparison of all cases and controls**

Summarizing the overall severity of mitochondrial impairment in the five disorders (**Figure 3**) indicates that major psychiatric disorders MDD, BD and SZ show less robust effects. The small effect sizes in those pathologies were most likely due to the small number of studies with small sample sizes, large region-specific heterogeneity and variation in medication and treatment, which made it harder to detect underlying mitochondrial disease. By contrast, strong evidence for a complex I deficit is found in PD, whereas a complex IV deficit is present in AD; with the simultaneous decline in complex I and IV being significant in both AD and PD. These results indicate an interaction of complex I with IV involved in the assembly and stability of complex I (Diaz *et al*, 2006; Li *et al*, 2007; Schäfer *et al*, 2007).

The interpretation of the strengths of the effect sizes between the five disorder compared with normal aging, as shown in **Figure 3**, should however be cautious. Aging is a major risk factor for developing both AD (Swerdlow, 2011) and idiopathic PD (Reeve *et al*, 2014).

Consequently, the effects of normal aging and that of the diseases cannot be clearly separated because of their inter-dependence. Of note, patients in all studies included were age-matched, thus in principle presenting age-adjusted effects. We therefore argue that the lesser effects in mood disorders is not related to the younger age of these patients, and the larger effects in AD is also not related to older age.

Further research is required to enlarge sample sizes and strengthen the evidence base. Our results may stimulate the application of novel in-vivo technologies such as assessing complex I NAD<sup>+</sup>/NADH redox state via auto-fluorescence (Blacker and Duchon, 2016) or magnetic resonance spectroscopy (Zhu *et al*, 2015) and complex IV redox states via functional near-infrared spectroscopy (Bale *et al*, 2016).

## **Supplementary materials**

**Datasets 1-6:** Datasets collected for the disorders MDD, BD, SZ, AD, PD and AGING.

## **Funding and Disclosure**

The work was funded by a Janssen Fellowship in Translational Neuroscience at Columbia University, New York awarded to LH.

Dr. Mann receives royalties for commercial use of the C-SSRS from the Research Foundation of Mental Hygiene. The other authors report no conflict of interest.

## **Acknowledgements**

The authors thank Tadafumi Kato, Brain Science Institute, RIKEN, Wako, Saitama, Japan, for sharing data. The authors thank Mercedes Armand-Ugon, Bellvitge University Hospital, Barcelona, Spain, for sharing data. The authors thank Tony Altar, Verge Genomics, San Francisco, USA, for sharing data.



## References

- Aarsland D, Cummings J, Larsen J (2001). Neuropsychiatric differences between Parkinson's disease with dementia and Alzheimer's disease. *Int J Geriatr Psychiatry* **16**: 184–191.
- Abreu R de S, Penalva LO, Marcotte EM, Vogel C (2009). Global signatures of protein and mRNA expression levels. *Molecular bioSystems* **5**: 1512–1526.
- Akarsu S, Torun D, Erdem M, Kozan S, Akar H, Uzun O (2015). Mitochondrial complex I and III mRNA levels in bipolar disorder. *Journal of Affective Disorders* **184**: 160–163.
- Aksenov MY, Tucker HM, Nair P, Aksenova MV, Butterfield DA, Estus S, *et al* (1999). The Expression of Several Mitochondrial and Nuclear Genes Encoding the Subunits of Electron Transport Chain Enzyme Complexes, Cytochrome c Oxidase, and NADH Dehydrogenase, in Different Brain Regions in Alzheimer's Disease. *Neurochemical Research* **24**: 767–774.
- Alikhani N, Guo L, Yan S, Du H, Pinho CM, Chen JX, *et al* (2011). Decreased Proteolytic Activity of the Mitochondrial Amyloid- $\beta$  Degrading Enzyme, PreP Peptidasome, in Alzheimer's Disease Brain Mitochondria. *Journal of Alzheimer's Disease* **27**: 75–87.
- Altar CA, Jurata LW, Charles V, Lemire A, Liu P, Bukhman Y, *et al* (2005). Deficient Hippocampal Neuron Expression of Proteasome, Ubiquitin, and Mitochondrial Genes in Multiple Schizophrenia Cohorts. *Biological Psychiatry* **58**: 85–96.
- American Psychiatric Association (American Psychiatric Press: Washington, DC, 2000). *Diagnostic and statistical manual of mental disorders (DSM-IV-TR)*. .
- Anderson JJ, Bravi D, Ferrari R, Davis TL, Baronti F, Chase TN, *et al* (1993). No evidence for altered muscle mitochondrial function in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry* **56**: 477–480.
- Andreazza AC, Shao L, Wang J, Young L (2010). Mitochondrial complex i activity and oxidative damage to mitochondrial proteins in the prefrontal cortex of patients with bipolar disorder. *Archives of General Psychiatry* **67**: 360–368.
- Andreazza AC, Wang J-F, Salmasi F, Shao L, Young LT (2013). Specific subcellular changes in oxidative stress in prefrontal cortex from patients with bipolar disorder. *J Neurochem* **127**: 552–561.
- Armand-Ugon M, Ansoleaga B, Berjaoui S, Ferrer I (2017). Reduced Mitochondrial Activity is Early and Steady in the Entorhinal Cortex but it is Mainly Unmodified in the Frontal Cortex in Alzheimer's Disease. *Current Alzheimer Research* **14**: 1327–1334.
- Arnold S (2012). Cytochrome c Oxidase and Its Role in Neurodegeneration and Neuroprotection. *Mitochondrial Oxidative Phosphorylation: Nuclear-Encoded Genes, Enzyme Regulation, and Pathophysiology* 305–339doi:10.1007/978-1-4614-3573-0\_13.
- Atkinson-Clement C, Pinto S, Eusebio A, Coulon O (2017). Diffusion tensor imaging in Parkinson's disease: Review and meta-analysis. *NeuroImage : Clinical* **16**: 98–110.
- Bale G, Elwell C, Tachtsidis I (2016). From Jöbsis to the Present Day: A Review of Clinical Near-Infrared Spectroscopy Measurements of Cerebral Cytochrome-C-Oxidase. *Journal of Biomedical Optics* **21**: 91307.
- Bansal Y, Kuhad A (2016). Mitochondrial Dysfunction in Depression. *Current Neuropharmacology* **14**: 610–618.
- Barroso N, Campos Y, Huertas R, Esteban J, Molina JA, Alonso A, *et al* (1993). Respiratory chain enzyme activities in lymphocytes from untreated patients with Parkinson disease. *Clin Chem* **39**: 667.
- Beech RD, Lowthert L, Leffert JJ, Mason PN, Taylor MM, Umlauf S, *et al* (2010). Increased peripheral blood expression of electron transport chain genes in bipolar depression. *Bipolar disorders* **12**: 813–824.
- Benecke R, Struemper P, Weiss H (1993). Electron transfer complexes I and IV of platelets are abnormal in Parkinson's disease but normal in Parkinson-plus syndromes. *Brain* **116**: 1451–63.
- Ben-Shachar D, Bonne O, Chisin R, Klein E, Lester H, Aharon-Peretz J, *et al* (2007). Cerebral glucose utilization and platelet mitochondrial complex I activity in schizophrenia: A FDG-PET study. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **31**: 807–813.



- Ben-Shachar D, Karry R (2007). Sp1 Expression Is Disrupted in Schizophrenia; A Possible Mechanism for the Abnormal Expression of Mitochondrial Complex I Genes, NDUFV1 and NDUFV2. *PLoS ONE* **2**: e817.
- Ben-Shachar D, Karry R (2008). Neuroanatomical Pattern of Mitochondrial Complex I Pathology Varies between Schizophrenia, Bipolar Disorder and Major Depression. *PLoS ONE* **3**: e3676.
- Ben-Shachar D, Zuk R, Gazawi H, Reshef A, Sheinkman A, Klein E (1999). Increased mitochondrial complex I activity in platelets of schizophrenic patients. *International Journal of Neuropsychopharmacology* **2**: 245–253.
- Bergman O, Ben-Shachar D (2016). Mitochondrial Oxidative Phosphorylation System (OXPHOS) Deficits in Schizophrenia: Possible Interactions with Cellular Processes. *Can J Psychiatry* **61**: 457–469.
- Bindoff LA, Birch-Machin MA, Cartledge NEF, Parker WD, Turnbull DM (1991). Respiratory chain abnormalities in skeletal muscle from patients with Parkinson's disease. *Journal of the Neurological Sciences* **104**: 203–208.
- Blacker TS, Duchon MR (2016). Investigating mitochondrial redox state using NADH and NADPH autofluorescence. *Free Radical Biology & Medicine* **100**: 53–65.
- Blake CI, Spitz E, Leehey M, Hoffer BJ, Boyson SJ (1997). Platelet mitochondrial respiratory chain function in Parkinson's disease. *Mov Disord* **12**: 3–8.
- Blin O, Desnuelle C, Rascol O, Borg M, Paul HPS, Azulay JP, *et al* (1994). Mitochondrial respiratory failure in skeletal muscle from patients with Parkinson's disease and multiple system atrophy. *Journal of the Neurological Sciences* **125**: 95–101.
- Boffoli D, Scacco SC, Vergari R, Persio MT, Solarino G, Laforgia R, *et al* (1996). Ageing is associated in females with a decline in the content and activity of the b-c1 complex in skeletal muscle mitochondria. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1315**: 66–72.
- Boffoli D, Scacco SC, Vergari R, Solarino G, Santacroce G, Papa S (1994). Decline with age of the respiratory chain activity in human skeletal muscle. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1226**: 73–82.
- Bonawitz ND, Clayton DA, Shadel GS (2006). Initiation and Beyond: Multiple Functions of the Human Mitochondrial Transcription Machinery. *Molecular Cell* **24**: 813–825.
- Bosetti F, Brizzi F, Barogi S, Mancuso M, Siciliano G, Tendi EA, *et al* (2002). Cytochrome c oxidase and mitochondrial F1F0-ATPase (ATP synthase) activities in platelets and brain from patients with Alzheimer's disease. *Neurobiology of Aging* **23**: 371–376.
- Boublay N, Schott AM, Krolak-Salmon P (2016). Neuroimaging correlates of neuropsychiatric symptoms in Alzheimer's disease: a review of 20 years of research. *Eur J Neurol* **23**: 1500–1509.
- Boveris A, Costa L, Cadenas E (1999). The mitochondrial production of oxygen radicals and cellular aging. *OXIDATIVE STRESS AND DISEASE* **2**: 1–16.
- Brakowski J, Spinelli S, Dörig N, Bosch OG, Manoliu A, Holtforth MG, *et al* (2017). Resting state brain network function in major depression – Depression symptomatology, antidepressant treatment effects, future research. *Journal of Psychiatric Research* **92**: 147–159.
- Bravi D, Anderson JJ, Dagani F, Davis TL, Ferrari R, Gillespie M, *et al* (1992). Effect of aging and dopaminomimetic therapy on mitochondrial respiratory function in Parkinson's disease. *Mov Disord* **7**: 228–231.
- Brisch R, Saniotis A, Wolf R, Biellau H, Bernstein H-G, Steiner J, *et al* (2014). The Role of Dopamine in Schizophrenia from a Neurobiological and Evolutionary Perspective: Old Fashioned, but Still in Vogue. *Frontiers in Psychiatry* **5**: 47.
- Bronstein JM, Paul K, Yang L, Haas RH, Shults CW, Le T, *et al* (2015). Platelet Mitochondrial Activity and Pesticide Exposure in Early Parkinson's Disease. *Movement disorders : official journal of the Movement Disorder Society* **30**: 862–866.
- Cabré R, Naudí A, Dominguez-Gonzalez M, Ayala V, Jové M, Mota-Martorell N, *et al* (2017). Sixty years old is the breakpoint of human frontal cortex aging. *Free Radical Biology and Medicine* **103**: 14–22.

- Cardellach F, Martí MJ, Fernández-Solá J, Marín C, Hoek JB, Tolosa E, *et al* (1993). Mitochondrial respiratory chain activity in skeletal muscle from patients with Parkinson's disease. *Neurology* **43**: 2258.
- Cardoso SM, Proença MT, Santos S, Santana I, Oliveira CR (2004). Cytochrome c oxidase is decreased in Alzheimer's disease platelets. *Neurobiology of Aging* **25**: 105–110.
- Casademont J, Rodriguez-Santiago B, Miró O, Beato A, López S, Nunes V, *et al* (2005). Mitochondrial respiratory chain in brain homogenates: activities in different brain areas in patients with Alzheimer's disease. *Aging Clinical and Experimental Research* **17**: 1–7.
- Cavelier L, Jazin EE, Eriksson I, Prince J, Båve U, Orelund L, *et al* (1995). Decreased Cytochrome-c Oxidase Activity and Lack of Age-Related Accumulation of Mitochondrial DNA Deletions in the Brains of Schizophrenics. *Genomics* **29**: 217–224.
- Cenini G, Rüb C, Bruderek M, Voos W, Gilmore R (2016). Amyloid  $\beta$ -peptides interfere with mitochondrial preprotein import competence by a coaggregation process. *MBoC* **27**: 3257–3272.
- Chagnon P, Bétard C, Robitaille Y, Cholette A, Gauvrea D (1995). Distribution of brain cytochrome oxidase activity in various neurodegenerative diseases. *Neuroreport: An International Journal for the Rapid Communication of Research in Neuroscience* **6**: 711–715.
- Chandrasekaran K, Giordano T, Brady DR, Stoll J, Martin LJ, Rapoport SI (1994). Impairment in mitochondrial cytochrome oxidase gene expression in Alzheimer disease. *Molecular Brain Research* **24**: 336–340.
- Chandrasekaran K, Hatanpää K, Brady DR, Stoll J, Rapoport SI (1998). Downregulation of oxidative phosphorylation in Alzheimer disease: loss of cytochrome oxidase subunit mRNA in the hippocampus and entorhinal cortex. *Brain Research* **796**: 13–19.
- Christian BE, Shadel GS (2014). Aging: It's SIRTainly Possible to Restore Mitochondrial Dysfunction. *Current Biology* **24**: R206–R208.
- Cooper JM, Daniel SE, Marsden CD, Schapira AHV (1995). L-Dihydroxyphenylalanine and complex I deficiency in Parkinson's disease brain. *Movement Disorders* **10**: 295–297.
- Cooper JM, Wischik C, Schapira AHV (1993). Mitochondrial function in Alzheimer's disease. *The Lancet* **341**: 969–970.
- Coskun P, Wyrembak J, Schriener S, Chen H-W, Marciniack C, LaFerla F, *et al* (2012). A Mitochondrial Etiology of Alzheimer and Parkinson Disease. *Biochimica et Biophysica Acta* **1820**: 553–564.
- Cottrell DA, Blakely EL, Johnson MA, Ince PG, Turnbull DM (2001). Mitochondrial enzyme-deficient hippocampal neurons and choroidal cells in AD. *Neurology* **57**: 260.
- Devi L, Prabhu BM, Galati DF, Avadhani NG, Anandatheerthavarada HK (2006). Accumulation of Amyloid Precursor Protein in the Mitochondrial Import Channels of Human Alzheimer's Disease Brain Is Associated with Mitochondrial Dysfunction. *J Neurosci* **26**: 9057.
- Diaz F, Fukui H, Garcia S, Moraes CT (2006). Cytochrome c Oxidase Is Required for the Assembly/Stability of Respiratory Complex I in Mouse Fibroblasts. *Molecular and Cellular Biology* **26**: 4872–4881.
- DiDonato S, Zeviani M, Giovannini P, Savarese N, Rimoldi M, Mariotti C, *et al* (1993). Respiratory chain and mitochondrial DNA in muscle and brain in Parkinson's disease patients. *Neurology* **43**: 2262.
- Dietsche B, Kircher T, Falkenberg I (2017). Structural brain changes in schizophrenia at different stages of the illness: A selective review of longitudinal magnetic resonance imaging studies. *Aust N Z J Psychiatry* **51**: 500–508.
- Distefano G, Standley RA, Dubé JJ, Carnero EA, Ritov VB, Stefanovic-Racic M, *et al* (2017). Chronological Age Does not Influence Ex-vivo Mitochondrial Respiration and Quality Control in Skeletal Muscle. *The Journals of Gerontology: Series A* **72**: 535–542.
- Dixit A, Srivastava G, Verma D, Mishra M, Singh PK, Prakash O, *et al* (2013). Minocycline, levodopa and MnTMPyP induced changes in the mitochondrial proteome profile of MPTP and maneb and paraquat mice models of Parkinson's disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1832**: 1227–1240.

- Dror N, Klein E, Karry R, Sheinkman A, Kirsh Z, Mazor M, *et al* (2002). State-dependent alterations in mitochondrial complex I activity in platelets: a potential peripheral marker for schizophrenia. *Molecular Psychiatry* **7**: 995.
- Duke DC, Moran LB, Pearce RKB, Graeber MB (2007). The medial and lateral substantia nigra in Parkinson's disease: mRNA profiles associated with higher brain tissue vulnerability. *Neurogenetics* **8**: 83–94.
- Emelyanova L, Preston C, Gupta A, Viqar M, Negmadjanov U, Edwards S, *et al* (2017). Effect of Aging on Mitochondrial Energetics in the Human Atria. *The Journals of Gerontology: Series A* glx160–glx160doi:10.1093/gerona/glx160.
- Feldhaus P, Fraga DB, Ghedim FV, De Luca RD, Bruna TD, Heluany M, *et al* (2011). Evaluation of respiratory chain activity in lymphocytes of patients with Alzheimer disease. *Metabolic Brain Disease* **26**: 229.
- Fišar Z, Hroudová J, Hansíková H, Spáčilová J, Lelková P, Wenchich L, *et al* (2016). Mitochondrial Respiration in the Platelets of Patients with Alzheimer's Disease. *Current Alzheimer Research* **13**: 930–941.
- Fukuyama R, Hatanpää K, Rapoport SI, Chandrasekaran K (1996). Gene expression of ND4, a subunit of complex I of oxidative phosphorylation in mitochondria, is decreased in temporal cortex of brains of Alzheimer's disease patients. *Brain Research* **713**: 290–293.
- Gandal MJ, Haney JR, Parikshak NN, Leppa V, Ramaswami G, Hartl C, *et al* (2018). Shared molecular neuropathology across major psychiatric disorders parallels polygenic overlap. *Science* **359**: 693.
- Gao L, Wu T (2016). The study of brain functional connectivity in Parkinson's disease. *Translational Neurodegeneration* **5**: 18.
- Gardner A, Johansson A, Wibom R, Nennesmo I, Döbeln U von, Hagenfeldt L, *et al* (2003). Alterations of mitochondrial function and correlations with personality traits in selected major depressive disorder patients. *Journal of Affective Disorders* **76**: 55–68.
- Gardner A, Salmaso D, Nardo D, Micucci F, Nobili F, Sanchez-Crespo A, *et al* (2008). Mitochondrial Function Is Related to Alterations at Brain SPECT in Depressed Patients. *CNS Spectrums* **13**: 805–814.
- Gatt AP, Duncan OF, Attems J, Francis PT, Ballard CG, Bateman JM (2016). Dementia in Parkinson's disease is associated with enhanced mitochondrial complex I deficiency. *Mov Disord* **31**: 352–359.
- Giachin G, Bouverot R, Acajjaoui S, Pantalone S, Soler-López M (2016). Dynamics of Human Mitochondrial Complex I Assembly: Implications for Neurodegenerative Diseases. *Frontiers in Molecular Biosciences* **3**: 43.
- Grossberg GT (2003). Cholinesterase Inhibitors for the Treatment of Alzheimer's Disease:: Getting On and Staying On. *Current Therapeutic Research, Clinical and Experimental* **64**: 216–235.
- Gu M, Owen A., Toffa SE., Cooper J., Dexter D., Jenner P, *et al* (1998). Mitochondrial function, GSH and iron in neurodegeneration and Lewy body diseases. *Journal of the Neurological Sciences* **158**: 24–29.
- Gubert C, Stertz L, Pfaffenseller B, Panizzutti BS, Rezin GT, Massuda R, *et al* (2013). Mitochondrial activity and oxidative stress markers in peripheral blood mononuclear cells of patients with bipolar disorder, schizophrenia, and healthy subjects. *Journal of Psychiatric Research* **47**: 1396–1402.
- Gueugneau M, Coudy-Gandilhon C, Théron L, Meunier B, Barboiron C, Combaret L, *et al* (2015). Skeletal Muscle Lipid Content and Oxidative Activity in Relation to Muscle Fiber Type in Aging and Metabolic Syndrome. *The Journals of Gerontology: Series A* **70**: 566–576.
- Haas RH, Nasirian F, Nakano K, Ward D, Pay M, Hill R, *et al* (1995). Low platelet mitochondrial complex I and complex II/III activity in early untreated parkinson's disease. *Ann Neurol* **37**: 714–722.
- Hanagasi HA, Ayribas D, Baysal K, Emre M (2005). Mitochondrial complex I, II/III, and IV activities in familial and sporadic Parkinson's disease. *International Journal of Neuroscience* **115**: 479–493.
- Harman D (1992). Free radical theory of aging. *Mutation Research/DNAging* **275**: 257–266.

- Hibar DP, Westlye LT, Doan NT, Jahanshad N, Cheung JW, Ching CRK, *et al* (2017). Cortical abnormalities in bipolar disorder: an MRI analysis of 6503 individuals from the ENIGMA Bipolar Disorder Working Group. *Molecular Psychiatry* at <<http://dx.doi.org/10.1038/mp.2017.73>>.
- Hirai K, Aliev G, Nunomura A, Fujioka H, Russell RL, Atwood CS, *et al* (2001). Mitochondrial abnormalities in Alzheimer's disease. *Journal of Neuroscience* **21**: 3017–3023.
- Hroudová J, Fišar Z (2011). Connectivity between mitochondrial functions and psychiatric disorders. *Psychiatry Clin Neurosci* **65**: 130–141.
- Hroudová J, Fišar Z, Korábečný J, Nepovimová E, Spilovská K, Vašková L, *et al* (2017). In vitro effects of newly developed cholinesterase inhibitors and ABAD modulators on mitochondrial respiration. *European Neuropsychopharmacology* **27**: S1022.
- Hsieh R, Hou J, Hsu H, Wei Y (1994). Age-dependent respiratory function decline and DNA deletions in human muscle mitochondria. *Biochemistry and Molecular Biology International* **32**: 10009–22.
- Iwamoto K, Bundo M, Kato T (2005). Altered expression of mitochondria-related genes in postmortem brains of patients with bipolar disorder or schizophrenia, as revealed by large-scale DNA microarray analysis. *Human Molecular Genetics* **14**: 241–253.
- Jiang J, Zhao Y-J, Hu X-Y, Du M-Y, Chen Z-Q, Wu M, *et al* (2017). Microstructural brain abnormalities in medication-free patients with major depressive disorder: a systematic review and meta-analysis of diffusion tensor imaging. *Journal of Psychiatry & Neuroscience : JPN* **42**: 150–163.
- Karry R, Klein E, Ben Shachar D (2004). Mitochondrial complex i subunits expression is altered in schizophrenia: a postmortem study. *Biological Psychiatry* **55**: 676–684.
- Kato T (2017). Neurobiological basis of bipolar disorder: Mitochondrial dysfunction hypothesis and beyond. *Schizophrenia Research* **187**: 62–66.
- Keeney PM, Xie J, Capaldi RA, Bennett JP (2006). Parkinson's Disease Brain Mitochondrial Complex I Has Oxidatively Damaged Subunits and Is Functionally Impaired and Misassembled. *J Neurosci* **26**: 5256.
- Kim HK, Andreazza AC, Elmi N, Chen W, Young LT (2016). Nod-like receptor pyrin containing 3 (NLRP3) in the post-mortem frontal cortex from patients with bipolar disorder: A potential mediator between mitochondria and immune-activation. *Journal of Psychiatric Research* **72**: 43–50.
- Kim SH, Vlkolinsky R, Cairns N, Fountoulakis M, Lubec G (2001). The reduction of NADH: Ubiquinone oxidoreductase 24- and 75-kDa subunits in brains of patients with Down syndrome and Alzheimer's disease. *Life Sciences* **68**: 2741–2750.
- Kish SJ, Bergeron C, Rajput A, Dozic S, Mastrogiamo F, Chang L-J, *et al* (1992). Brain Cytochrome Oxidase in Alzheimer's Disease. *Journal of Neurochemistry* **59**: 776–779.
- Kish SJ, Mastrogiamo F, Guttman M, Furukawa Y, Taanman J-W, Dozic S, *et al* (1999). Decreased Brain Protein Levels of Cytochrome Oxidase Subunits in Alzheimer's Disease and in Hereditary Spinocerebella Ataxia Disorders. *Journal of Neurochemistry* **72**: 700–707.
- Konradi C, Eaton M, MacDonald ML, Walsh J, Benes FM, Heckers S (2004). Molecular evidence for mitochondrial dysfunction in bipolar disorder. *Archives of General Psychiatry* **61**: 300–308.
- Korábečný J, Nepovimová E, Cikánková T, Špilovská K, Vašková L, Mezeiová E, *et al* (2018). Newly Developed Drugs for Alzheimer's Disease in Relation to Energy Metabolism, Cholinergic and Monoaminergic Neurotransmission. *Neuroscience* **370**: 191–206.
- Krige D, Carroll M, Cooper J, Marsden C, Schapira A (1992). Platelet mitochondrial function in Parkinson's disease. *Annals of Neurology* **32**: 782–8.
- Lancôt KL, Amatniek J, Ancoli-Israel S, Arnold SE, Ballard C, Cohen-Mansfield J, *et al* (2017). Neuropsychiatric signs and symptoms of Alzheimer's disease: New treatment paradigms. *Alzheimer's & Dementia : Translational Research & Clinical Interventions* **3**: 440–449.
- Lanza IR, Short DK, Short KR, Raghavakaimal S, Basu R, Joyner MJ, *et al* (2008). Endurance Exercise as a Countermeasure for Aging. *Diabetes* **57**: 2933–2942.
- Li Y, D'Aurelio M, Deng J-H, Park J-S, Manfredi G, Hu P, *et al* (2007). An Assembled Complex IV Maintains the Stability and Activity of Complex I in Mammalian Mitochondria. *Journal of Biological Chemistry* **282**: 17557–17562.

- Liang WS, Reiman EM, Valla J, Dunckley T, Beach TG, Grover A, *et al* (2008). Alzheimer's disease is associated with reduced expression of energy metabolism genes in posterior cingulate neurons. *Proceedings of the National Academy of Sciences of the United States of America* **105**: 4441–4446.
- Long J, He P, Shen Y, Li R (2012). New evidence of mitochondria dysfunction in the female Alzheimer's brain: deficiency of estrogen receptor- $\beta$ . *Journal of Alzheimer's disease: JAD* **30**: 545–558.
- Lunnon K, Ibrahim Z, Proitsi P, Lourdasamy A, Newhouse S, Sattlecker M, *et al* (2012). Mitochondrial Dysfunction and Immune Activation are Detectable in Early Alzheimer's Disease Blood. *Journal of Alzheimer's Disease* **30**: 685–710.
- Lyketsos CG, Peters ME (2015). Dementia in patients with schizophrenia: Evidence for heterogeneity. *JAMA Psychiatry* **72**: 1075–1076.
- Maier T, Güell M, Serrano L (2009). Correlation of mRNA and protein in complex biological samples. *FEBS Letters* **583**: 3966–3973.
- Mamah D, Wang L, Erausquin DBGA de, Gado M, Csernansky JG (2007). Structural Analysis of the Basal Ganglia in Schizophrenia. *Schizophrenia research* **89**: 59–71.
- Mancuso M, Filosto M, Bosetti F, Ceravolo R, Rocchi A, Tognoni G, *et al* (2003). Decreased platelet cytochrome c oxidase activity is accompanied by increased blood lactate concentration during exercise in patients with Alzheimer disease. *Experimental Neurology* **182**: 421–426.
- Manczak M, Park BS, Jung Y, Reddy PH (2004). Differential expression of oxidative phosphorylation genes in patients with Alzheimer's disease. *NeuroMolecular Medicine* **5**: 147–162.
- Mandavilli BS, Santos JH, Van Houten B (2002). Mitochondrial DNA repair and aging. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **509**: 127–151.
- Manji H, Kato T, Di Prospero NA, Ness S, Beal MF, Krams M, *et al* (2012). Impaired mitochondrial function in psychiatric disorders. *Nature Reviews Neuroscience* **13**: 293.
- Mann VM, Cooper JM, Krige D, Daniel SE, Schapira AHV, Marsden CD (1992). Brain, skeletal muscle and platelet homogenate mitochondrial function in Parkinson's disease. *Brain* **115**: 333–342.
- Martín MA, Molina JA, Jiménez-Jiménez FJ, Benito-León J, Ortí-Pareja M, Campos Y, *et al* (1996). Respiratory-chain enzyme activities in isolated mitochondria of lymphocytes from untreated Parkinson's disease patients. *Neurology* **46**: 1343.
- Maurer I, Zierz S, Möller H (2000). A selective defect of cytochrome c oxidase is present in brain of Alzheimer disease patients. *Neurobiology of Aging* **21**: 455–462.
- Maurer I, Zierz S, Möller H-J (2001). Evidence for a mitochondrial oxidative phosphorylation defect in brains from patients with schizophrenia. *Schizophrenia Research* **48**: 125–136.
- Mehler-Wex C, Duvigneau JC, Hartl RT, Ben-Shachar D, Warnke A, Gerlach M (2006). Increased mRNA levels of the mitochondrial complex I 75-kDa subunit. *European Child & Adolescent Psychiatry* **15**: 504–507.
- Merlo Pich M, Bovina C, Formiggini G, Cometti GG, Ghelli A, Parenti Castelli G, *et al* (1996). Inhibitor sensitivity of respiratory complex I in human platelets: A possible biomarker of ageing. *FEBS Letters* **380**: 176–178.
- Merlo Pich M, Raule N, Catani L, Fagioli ME, Faenza I, Cocco L, *et al* (2004). Increased transcription of mitochondrial genes for Complex I in human platelets during ageing. *FEBS Letters* **558**: 19–22.
- Miquel J, Economos AC, Fleming J, Johnson JE (1980). Mitochondrial role in cell aging. *Experimental Gerontology* **15**: 575–591.
- Mizuno Y, Suzuki K, Ohta S (1990). Postmortem changes in mitochondrial respiratory enzymes in brain and a preliminary observation in Parkinson's disease. *Journal of the Neurological Sciences* **96**: 49–57.
- Munkholm K, Peijs L, Vinberg M, Kessing LV (2015). A composite peripheral blood gene expression measure as a potential diagnostic biomarker in bipolar disorder. *Translational Psychiatry* **5**: e614.
- Mutisya EM, Bowling AC, Beal MF (1994). Cortical Cytochrome Oxidase Activity Is Reduced in Alzheimer's Disease. *Journal of Neurochemistry* **63**: 2179–2184.

- Mythri RB, Venkateshappa C, Harish G, Mahadevan A, Muthane UB, Yasha TC, *et al* (2011). Evaluation of Markers of Oxidative Stress, Antioxidant Function and Astrocytic Proliferation in the Striatum and Frontal Cortex of Parkinson's Disease Brains. *Neurochemical Research* **36**: 1452–1463.
- Nakagawa-Hattori Y, Yoshino H, Kondo T, Mizuno Y, Horai S (1992). Is Parkinson's disease a mitochondrial disorder? *Journal of the Neurological Sciences* **107**: 29–33.
- Navarro A, Boveris A (2004). Rat brain and liver mitochondria develop oxidative stress and lose enzymatic activities on aging. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **287**: R1244–R1249.
- Navarro A, Boveris A (2007). The mitochondrial energy transduction system and the aging process. *Am J Physiol Cell Physiol* **292**: C670.
- Naydenov AV, MacDonald ML, Ongur D, Konradi C (2007). Differences in lymphocyte electron transport gene expression levels between subjects with bipolar disorder and normal controls in response to glucose deprivation stress. *Archives of General Psychiatry* **64**: 555–564.
- Ogborn DI, McKay BR, Crane JD, Safdar A, Akhtar M, Parise G, *et al* (2015). Effects of age and unaccustomed resistance exercise on mitochondrial transcript and protein abundance in skeletal muscle of men. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **308**: R734–R741.
- Ojaimi J, Masters CL, Opeskin K, McKelvie P, Byrne E (1999). Mitochondrial respiratory chain activity in the human brain as a function of age. *Mechanisms of Ageing and Development* **111**: 39–47.
- Onyango I, Khan S, Bennett J (2017). Mitochondria in the pathophysiology of Alzheimer's and Parkinson's diseases. *Frontiers in Bioscience* **22**: 854–872.
- Pacifico R, Davis RL (2016). Transcriptome sequencing implicates dorsal striatum-specific gene network, immune response and energy metabolism pathways in bipolar disorder. *Molecular Psychiatry* **22**: 441.
- Parker W, Boyson S, Parks J (1989). Abnormalities of the electron transport chain in idiopathic Parkinson's disease. *Annals of Neurology* **26**: 719–23.
- Parker W, Filley C, Parks J (1990). Cytochrome oxidase deficiency in Alzheimer's disease. *Neurology* **40**: 1302–3.
- Parker W, Mahr N, Filley C, Parks J, Hughes D, Young D, *et al* (1994a). Reduced platelet cytochrome c oxidase activity in Alzheimer's disease. *Neurology* **44**: 1086–1090.
- Parker W, Parks J, Filley C, Kleinschmidt-DeMasters B (1994b). Electron transport chain defects in Alzheimer's disease brain. *Neurology* **44**: 1090–96.
- Parker WD, Parks JK, Swerdlow RH (2008). Complex I deficiency in Parkinson's disease frontal cortex. *Brain Research* **1189**: 215–218.
- Payne B, Chinnery P (2015). Mitochondrial dysfunction in aging: Much progress but many unresolved questions. *Biochimica et Biophysica Acta* **1847**: 1347–53.
- Pérez-Gracia E, Torrejón-Escribano B, Ferrer I (2008). Dystrophic neurites of senile plaques in Alzheimer's disease are deficient in cytochrome c oxidase. *Acta Neuropathologica* **116**: 261–268.
- Perluigi M, Barone E, Di Domenico F, Butterfield DA (2016). Aberrant protein phosphorylation in Alzheimer disease brain disturbs pro-survival and cell death pathways. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1862**: 1871–1882.
- Pestronk A, Keeling R, Choksi R (2017). Sarcopenia, age, atrophy, and myopathy: Mitochondrial oxidative enzyme activities. *Muscle Nerve* **56**: 122–128.
- Picone P, Nuzzo D, Caruana L, Scafidi V, Di Carlo M (2014). Mitochondrial Dysfunction: Different Routes to Alzheimer's Disease Therapy. *Oxidative Medicine and Cellular Longevity* **7**: 179.
- Pinho CM, Teixeira PF, Glaser E (2014). Mitochondrial import and degradation of amyloid- $\beta$  peptide. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1837**: 1069–1074.
- Prince JA, Blennow K, Gottfries CG, Karlsson I, Oreland L (1999). Mitochondrial Function is Differentially Altered in the Basal Ganglia of Chronic Schizophrenics. *Neuropsychopharmacology* **21**: 372.

- Prince JA, Harro J, Blennow K, Gottfries CG, Oreland L (2000). Putamen Mitochondrial Energy Metabolism Is Highly Correlated to Emotional and Intellectual Impairment in Schizophrenics. *Neuropsychopharmacology* **22**: 284.
- R Development Core Team (2008). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. .
- Rasmussen UF, Krstrup P, Kjaer M, Rasmussen HN (2003). Human skeletal muscle mitochondrial metabolism in youth and senescence: no signs of functional changes in ATP formation and mitochondrial oxidative capacity. *Pflügers Archiv* **446**: 270–278.
- Readnower R, Sauerbeck A, Sullivan P (2011). Mitochondria, Amyloid  $\beta$ , and Alzheimer's Disease,. *International Journal of Alzheimer's Disease* **Article ID 104545**: .
- Reeve A, Simcox E, Turnbull D (2014). Ageing and Parkinson's disease: Why is advancing age the biggest risk factor?(). *Ageing Research Reviews* **14**: 19–30.
- Reichmann H, Fhirke S, Hebenstreit G, Schrubar H, Riederer P (1993). Analyses of energy metabolism and mitochondrial genome in post-mortem brain from patients with Alzheimer's disease. *Journal of Neurology* **240**: 377–380.
- Rice MW, Smith KL, Roberts RC, Perez-Costas E, Melendez-Ferro M (2014). Assessment of Cytochrome C Oxidase Dysfunction in the Substantia Nigra/Ventral Tegmental Area in Schizophrenia. *PLoS ONE* **9**: e100054.
- Rooyackers OE, Adey DB, Ades PA, Nair KS (1996). Effect of age on in vivo rates of mitochondrial protein synthesis in human skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 15364–15369.
- Rosenfeld M, Brenner-Lavie H, Ari SG-B, Kavushansky A, Ben-Shachar D (2011). Perturbation in Mitochondrial Network Dynamics and in Complex I Dependent Cellular Respiration in Schizophrenia. *Biological Psychiatry* **69**: 980–988.
- Ryan BJ, Hoek S, Fon EA, Wade-Martins R (2015). Mitochondrial dysfunction and mitophagy in Parkinson's: from familial to sporadic disease. *Trends in Biochemical Sciences* **40**: 200–210.
- Safdar A, Hamadeh MJ, Kaczor JJ, Raha S, deBeer J, Tarnopolsky MA (2010). Aberrant Mitochondrial Homeostasis in the Skeletal Muscle of Sedentary Older Adults. *PLOS ONE* **5**: e10778.
- Sanchez-Bahillo A, Bautista-Hernandez V, Barcia Gonzalez C, Bañon R, Luna A, Hirsch E, *et al* (2008). Increased mRNA expression of cytochrome oxidase in dorsal raphe nucleus of depressive suicide victims. *Neuropsychiatric Disease and Treatment* **4**: 413–416.
- Schäfer E, Dencher NA, Vonck J, Parcej DN (2007). Three-Dimensional Structure of the Respiratory Chain Supercomplex I1III2IV1 from Bovine Heart Mitochondria,. *Biochemistry* **46**: 12579–12585.
- Schägger H (1995). Quantification of oxidative phosphorylation enzymes after blue native electrophoresis and two-dimensional resolution: Normal complex I protein amounts in Parkinson's disease conflict with reduced catalytic activities. *ELECTROPHORESIS* **16**: 763–770.
- Schapira AHV, Cooper JM, Dexter D, Jenner P, Clark JB, Marsden CD (1990a). Mitochondrial Complex I Deficiency in Parkinson's Disease. *The Lancet* **333**: 1269.
- Schapira AHV, Mann VM, Cooper JM, Dexter D, Daniel SE, Jenner P, *et al* (1990b). Anatomic and Disease Specificity of NADH CoQ1 Reductase (Complex I) Deficiency in Parkinson's Disease. *Journal of Neurochemistry* **55**: 2142–2145.
- Sekar S, McDonald J, Cuyugan L, Aldrich J, Kurdoglu A, Adkins J, *et al* (2015). Alzheimer's disease is associated with altered expression of genes involved in immune response and mitochondrial processes in astrocytes. *Neurobiology of Aging* **36**: 583–591.
- Shao L, Martin MV, Watson SJ, Schatzberg A, Akil H, Myers RM, *et al* (2008). Mitochondrial involvement in psychiatric disorders. *Annals of medicine* **40**: 281–295.
- Sheehan JP, Swerdlow RH, Miller SW, Davis RE, Parks JK, Parker WD, *et al* (1997). Calcium Homeostasis and Reactive Oxygen Species Production in Cells Transformed by Mitochondria from Individuals with Sporadic Alzheimer's Disease. *J Neurosci* **17**: 4612.
- Shinde S, Pasupathy K (2006). Respiratory-chain enzyme activities in isolated mitochondria of lymphocytes from patients with Parkinson's disease: Preliminary study. *Neurology India* **54**: 390–3.

- Shinn AK, Roh YS, Ravichandran CT, Baker JT, Öngür D, Cohen BM (2017). Aberrant Cerebellar Connectivity in Bipolar Disorder With Psychosis. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **2**: 438–448.
- Shoffner JM, Watts RL, Juncos JL, Torroni A, Wallace DC (1991). Mitochondrial oxidative phosphorylation defects in parkinson's disease. *Ann Neurol* **30**: 332–339.
- Shults CW, Nasirian F, Ward DM, Nakano K, Pay M, Hill LR, *et al* (1995). Cardidopa/Levodopa and selegiline do not affect platelet mitochondrial function in early Parkinsonism. *Neurology* **45**: 344–8.
- Simonian N, Hyman B (1993). Functional alterations in Alzheimer's disease: diminution of cytochrome oxidase in the hippocampal formation. *Journal of Neuropathology and Experimental Neurology* **52**: 580–5.
- Simonian N, Hyman B (1995). Functional alterations in Alzheimer's disease: selective loss of mitochondrial-encoded cytochrome oxidase mRNA in the hippocampal formation. *Journal of Neuropathology and Experimental Neurology* **53**: 508–12.
- Sousa RT de, Streck EL, Zanetti MV, Ferreira GK, Diniz BS, Brunoni AR, *et al* (2015). Lithium increases leukocyte mitochondrial complex I activity in bipolar disorder during depressive episodes. *Psychopharmacology* **232**: 245–250.
- Srinivasan S, Avadhani NG (2012). Cytochrome c Oxidase Dysfunction in Oxidative Stress. *Free radical biology & medicine* **53**: 1252–1263.
- Srinivasan V, Cardinali DP, Srinivasan US, Kaur C, Brown GM, Spence DW, *et al* (2011). Therapeutic potential of melatonin and its analogs in Parkinson's disease: focus on sleep and neuroprotection. *Therapeutic Advances in Neurological Disorders* **4**: 297–317.
- Sun N, Youle RJ, Finkel T (2016). The Mitochondrial Basis of Aging. *Molecular cell* **61**: 654–666.
- Sun X, Wang J-F, Tseng M, Young LT (2006). Downregulation in components of the mitochondrial electron transport chain in the postmortem frontal cortex of subjects with bipolar disorder. *Journal of Psychiatry and Neuroscience* **31**: 189–196.
- Swerdlow RH (2011). Brain aging, Alzheimer's disease, and mitochondria. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1812**: 1630–1639.
- Taurines R, Thome J, Duvigneau JC, Forbes-Robertson S, Yang L, Klampfl K, *et al* (2010). Expression analyses of the mitochondrial complex I 75-kDa subunit in early onset schizophrenia and autism spectrum disorder: increased levels as a potential biomarker for early onset schizophrenia. *European Child & Adolescent Psychiatry* **19**: 441–448.
- Terni B, Boada J, Portero-Otin M, Pamplona R, Ferrer I (2010). Mitochondrial ATP-Synthase in the Entorhinal Cortex Is a Target of Oxidative Stress at Stages I/II of Alzheimer's Disease Pathology. *Brain Pathology* **20**: 222–233.
- Thomas R, Keeney P, Bennett J (2012). Impaired Complex-I Mitochondrial Biogenesis in Parkinson Disease Frontal Cortex. *Journal of Parkinsons Disease* **2**: 67–76.
- Trounce I, Byrne E, Marzuki S (1989). Decline in skeletal muscle mitochondrial respiratory chain function: possible factor in ageing. *The Lancet* **333**: 637–639.
- Valla J, Berndt JD, Gonzalez-Lima F (2001). Energy Hypometabolism in Posterior Cingulate Cortex of Alzheimer's Patients: Superficial Laminar Cytochrome Oxidase Associated with Disease Duration. *J Neurosci* **21**: 4923.
- Valla J, Schneider L, Niedzielko T, Coon KD, Caselli R, Sabbagh MN, *et al* (2006). Impaired Platelet Mitochondrial Activity in Alzheimer's Disease and Mild Cognitive Impairment. *Mitochondrion* **6**: 323–330.
- Varghese M, Pandey M, Samanta A, Gangopadhyay PK, Mohanakumar KP (2009). Reduced NADH coenzyme Q dehydrogenase activity in platelets of Parkinson's disease, but not Parkinson plus patients, from an Indian population. *Journal of the Neurological Sciences* **279**: 39–42.
- Vawter M, Tomita H, Meng F, Bolstad B, Li J, Evans S, *et al* (2006). Mitochondrial-related gene expression changes are sensitive to agonal-pH state: implications for brain disorders. *Molecular psychiatry* **11**: 615–679.
- Verwer RWH, Jansen KA, Sluiter AA, Pool CW, Kamphorst W, Swaab DF (2000). Decreased Hippocampal Metabolic Activity in Alzheimer Patients Is Not Reflected in the Immunoreactivity of Cytochrome Oxidase Subunits. *Experimental Neurology* **163**: 440–451.



- Viechtbauer W (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software* **36**: 1–48.
- Vitali M, Venturelli E, Galimberti D, Benerini Gatta L, Scarpini E, Finazzi D (2009). Analysis of the genes coding for subunit 10 and 15 of cytochrome c oxidase in Alzheimer's disease. *Journal of Neural Transmission* **116**: 1635–1641.
- Vogel C, Marcotte EM (2012). Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nature reviews Genetics* **13**: 227–232.
- Washizuka S, Iwamoto K, Kakiuchi C, Bundo M, Kato T (2009). Expression of mitochondrial complex I subunit gene NDUFV2 in the lymphoblastoid cells derived from patients with bipolar disorder and schizophrenia. *Neuroscience Research* **63**: 199–204.
- Washizuka S, Kakiuchi C, Mori K, Tajima O, Akiyama T, Kato T (2005). Expression of mitochondria-related genes in lymphoblastoid cells from patients with bipolar disorder. *Bipolar Disorders* **7**: 146–152.
- Whatley S, Curti D, Marchbanks R (1996). Mitochondrial involvement in schizophrenia and other functional psychoses. *Neurochemical Research* **21**: 995–1004.
- Wiedemann FR, Winkler K, Lins H, Wallesch C-W, Kunz WS (1999). Detection of Respiratory Chain Defects in Cultivated Skin Fibroblasts and Skeletal Muscle of Patients with Parkinson's Disease. *Annals of the New York Academy of Sciences* **893**: 426–429.
- Womer FY, Wang L, Alpert K, Smith MJ, Csernansky JG, Barch D, *et al* (2014). Basal ganglia and thalamic morphology in schizophrenia and bipolar disorder. *Psychiatry research* **223**: 75–83.
- Wong-Riley M, Antuono P, Ho K-C, Egan R, Hevner R, Liebl W, *et al* (1997). Cytochrome oxidase in Alzheimer's disease: Biochemical, histochemical, and immunohistochemical analyses of the visual and other systems. *Vision Research* **37**: 3593–3608.
- Yoshino H, Nakagawa-Hattori Y, Kondo T, Mizuno Y (1992). Mitochondrial complex I and II activities of lymphocytes and platelets in Parkinson's disease. *Journal of Neural Transmission - Parkinson's Disease and Dementia Section* **4**: 27–34.
- Zhu X-H, Lu M, Lee B-Y, Ugurbil K, Chen W (2015). In vivo NAD assay reveals the intracellular NAD contents and redox state in healthy human brain and their age dependences. *Proceedings of the National Academy of Sciences of the United States of America* **112**: 2876–2881.
- Zhuo C, Xue R, Luo L, Ji F, Tian H, Qu H, *et al* (2017). Efficacy of antidepressive medication for depression in Parkinson disease: a network meta-analysis. *Medicine* **96**: e6698.
- Zucchini C, Pugnali A, Pallotti F, Solmi R, Crimi M, Castaldini C, *et al* (1995). Human skeletal muscle mitochondria in aging: lack of detectable morphological and enzymic defects. *Biochemistry and Molecular Biology International* **37**: 607–16.

## Figure legends

**Figure 1. Forest plots MDD, BD and SZ.** Multivariate random-effect estimates of the SMD (95% CI, p-values) of complex I (NDU) and IV (COX) enzyme activity and subunits are shown both in numerical and graphical form. Values lower than 0 indicate that patients had lower levels than controls (CO), and vice versa for values greater than 0; the dashed vertical line at SMD = 0 indicates no effect. The size of the filled circles for each estimated SMD is proportional to the weight of the studies. P = putamen, NA = nucleus accumbens, GP = globus pallidus.

**Figure 2. Forest plots AD, PD and AGING.** Multivariate random-effect estimates of the SMD (95% CI, p-values) of complex I (NDU) and IV (COX) enzyme activity and subunits are shown both in numerical and graphical form. Values lower than 0 indicate that patients (or older (O) individuals) had lower levels than controls (CO) (or younger (Y) individuals), and vice versa for values greater than 0; the dashed vertical line at SMD = 0 indicates no effect. The size of the filled circles for each estimated SMD is proportional to the weight of the studies.

**Figure 3. Forest plot Summary.** Multivariate random-effect estimates of the SMD (95% CI, p-values) of complex I and IV are shown both in numerical and graphical form. SMD are ordered according to their strengths. Values lower than 0 indicate that patients (P) had lower levels than controls (CO), and vice versa for values greater than 0; the dashed vertical line at SMD = 0 indicates no effect. The size of the filled circles for each estimated SMD is proportional to the weight of the studies.

**Figure 4. Schematic overview of our findings.** Highlighted are the main findings of the meta-analyses for the five disorders MDD, BD, SZ, AD, PD and normal aging. Region-specific heterogeneity means that complex levels vary from brain region to brain region (i.e.,

variation from ROI to ROI); tissue-specific heterogeneity indicates that complex levels vary between tissues (i.e., peripheral blood versus brain).

## Tables


**Table 1. Eligible studies.** 125 studies were eligible for the meta-analyses, part of which was included in the meta-analyses for complex I and IV (marked in gray). Some studies reported more than one disorder. The detailed data extracted from these studies are provided in the supplementary materials (**Tables S1-6**).


MDD (N = 7)	BD (N = 19)	SZ (N = 21)	AD (N = 43)	PD (N = 35)	AGING (N = 18)
Andreazza 2010 (Andreazza AC <i>et al</i> , 2010)	Akarsu 2015 (Akarsu <i>et al</i> , 2015)	Altar 2005 (Altar <i>et al</i> , 2005)	Aksenov 1999 (Aksenov <i>et al</i> , 1999)	Anderson 1993 (Anderson <i>et al</i> , 1993)	Boffoli 1994 (Boffoli <i>et al</i> , 1994)
Ben-Shachar 1999 (Ben-Shachar <i>et al</i> , 1999)	Altar 2005 (Altar <i>et al</i> , 2005)	Andreazza 2010 (Andreazza AC <i>et al</i> , 2010)	Alikhani 2012 (Alikhani <i>et al</i> , 2011)	Barroso 1993 (Barroso <i>et al</i> , 1993)	Boffoli 1996 (Boffoli <i>et al</i> , 1996)
Ben-Shachar 2008 (Ben-Shachar and Karry, 2008)	Andreazza 2010 (Andreazza AC <i>et al</i> , 2010)	Andreazza 2013 (Andreazza <i>et al</i> , 2013)	Armand-Ugon 2017 (Armand-Ugon <i>et al</i> , 2017)	Benecke 1993 (Benecke <i>et al</i> , 1993)	Cabr� 2017 (Cabr� <i>et al</i> , 2017)
Gardner 2003 (Gardner <i>et al</i> , 2003)	Andreazza 2013 (Andreazza <i>et al</i> , 2013)	Ben-Shachar 1999 (Ben-Shachar <i>et al</i> , 1999)	Bosetti 2002 (Bosetti <i>et al</i> , 2002)	Bindoff 1991 (Bindoff <i>et al</i> , 1991)	Distefano 2017 (Distefano <i>et al</i> , 2017)
Gardner 2008 (Gardner <i>et al</i> , 2008)	Beech 2010 (Beech <i>et al</i> , 2010)	Ben-Shachar 2007 (Ben-Shachar <i>et al</i> , 2007)	Cardoso 2004 (Cardoso <i>et al</i> , 2004)	Blake 1997 (Blake <i>et al</i> , 1997)	Emelyanova 2017 (Emelyanova <i>et al</i> , 2017)
Karry 2004 (Karry <i>et al</i> , 2004)	Ben-Shachar 1999 (Ben-Shachar <i>et al</i> , 1999)	Ben-Shachar 2008 (Ben-Shachar and Karry, 2008)	Casademont 2005 (Casademont <i>et al</i> , 2005)	Blin 1994 (Blin <i>et al</i> , 1994)	Gueugneau 2014 (Gueugneau <i>et al</i> , 2015)
Sanchez-Bahillo 2008 (Sanchez-Bahillo <i>et al</i> , 2008)	Ben-Shachar 2008 (Ben-Shachar and Karry, 2008)	Cavelier 1995 (Cavelier <i>et al</i> , 1995)	Cavelier 1995 (Cavelier <i>et al</i> , 1995)	Bravi 1992 (Bravi <i>et al</i> , 1992)	Hsieh 1994 (Hsieh <i>et al</i> , 1994)
	de Sousa 2015 (de Sousa <i>et al</i> , 2015)	Dror 2002 (Dror <i>et al</i> , 2002)	Chagnon 1995 (Chagnon <i>et al</i> , 1995)	Bronstein 2015 (Bronstein <i>et al</i> , 2015)	Lanza 2008 (Lanza <i>et al</i> , 2008)
	Gubert 2013 (Gubert <i>et al</i> , 2013)	Gubert 2013 (Gubert <i>et al</i> , 2013)	Chandrasekaran 1994 (Chandrasekaran <i>et al</i> , 1994)	Cardellach 1993 (Cardellach <i>et al</i> , 1993)	Merlo Pich 2004 (Merlo Pich <i>et al</i> , 2004)
	Iwamoto 2005 (Iwamoto <i>et al</i> , 2005)	Iwamoto 2005 (Iwamoto <i>et al</i> , 2005)	Chandrasekaran 1998 (Chandrasekaran <i>et al</i> , 1998)	Chagnon 1995 (Chagnon <i>et al</i> , 1995)	Merlo Pich 1996 (Merlo Pich <i>et al</i> , 1996)
	Karry 2004 (Karry <i>et al</i> , 2004)	Karry 2004 (Karry <i>et al</i> , 2004)	Cooper 1993 (Cooper <i>et al</i> , 1993)	Cooper 1995 (Cooper <i>et al</i> , 1995)	Ogborn 2015 (Ogborn <i>et al</i> , 2015)
	Kim 2016 (Kim <i>et al</i> , 2016)	Kim 2016 (Kim <i>et al</i> , 2016)	Cottrell 2001 (Cottrell <i>et al</i> , 2001)	DiDonato 1993 (DiDonato <i>et al</i> , 1993)	Ojaimi 1999 (Ojaimi <i>et al</i> , 1999)
	Konradi 2004 (Konradi C <i>et al</i> , 2004)	Maurer 2001 (Maurer <i>et al</i> , 2001)	Devi 2006 (Devi <i>et al</i> , 2006)	Duke 2007 (Duke <i>et al</i> , 2007)	Pestronk 2016 (Pestronk <i>et al</i> , 2017)
	Munkholm 2015 (Munkholm <i>et al</i> , 2015)	Mehler-Wex 2006 (Mehler-Wex <i>et al</i> , 2006)	Feldhaus 2011 (Feldhaus <i>et al</i> , 2011)	Gatt 2016 (Gatt <i>et al</i> , 2016)	Rasmussen 2003 (Rasmussen <i>et al</i> , 2003)
	Naydenov 2007 (Naydenov AV <i>et al</i> , 2007)	Prince 1999 (Prince <i>et al</i> , 1999)	Fi�ar 2016 (Fi�ar <i>et al</i> , 2016)	Gu 1998 (Gu <i>et al</i> , 1998)	Rooyackers 1996 (Rooyackers <i>et al</i> , 1996)
	Rosenfeld 2011 (Rosenfeld <i>et al</i> , 2011)	Prince 2000 (Prince <i>et al</i> , 2000)	Fukuyama 1996 (Fukuyama <i>et al</i> , 1996)	Haas 1995 (Haas <i>et al</i> , 1995)	Safdar 2010 (Safdar <i>et al</i> , 2010)
	Sun 2006 (Sun <i>et al</i> , 2006)	Rice 2014 (Rice <i>et al</i> , 2014)	Gu 1998 (Gu <i>et al</i> , 1998)	Hanagasi 2004 (Hanagasi <i>et al</i> , 2005)	Trounce 1989 (Trounce <i>et al</i> , 1989)
	Washizuka 2005 (Washizuka <i>et al</i> , 2005)	Rosenfeld 2011 (Rosenfeld <i>et al</i> , 2011)	Hirai 2001 (Hirai <i>et al</i> , 2001)	Keeney 2006 (Keeney <i>et al</i> , 2006)	Zucchini 1995 (Zucchini <i>et al</i> , 1995)
	Washizuka 2009 (Washizuka <i>et al</i> , 2009)	Taurines 2010 (Taurines <i>et al</i> , 2010)	Kim 2001 (Kim <i>et al</i> , 2001)	Krige 1992 (Krige <i>et al</i> , 1992)	
		Washizuka 2009 (Washizuka <i>et al</i> , 2009)	Kish 1992 (Kish <i>et al</i> , 1992)	Mann 1992 (Mann <i>et al</i> , 1992)	
		Whatley 1998 (Whatley <i>et al</i> , 1996)	Kish 1999 (Kish <i>et al</i> , 1999)	Martin 1996 (Martin <i>et al</i> , 1996)	
			Liang 2008 (Liang <i>et al</i> , 2008)	Mizuno 1990 (Mizuno <i>et al</i> , 1990)	
			Long 2012 (Long <i>et al</i> , 2012)	Mythri 2011 (Mythri <i>et al</i> , 2011)	
			Lunnon 2012 (Lunnon <i>et al</i> , 2012)	Nakagawa-Hattori 1992 (Nakagawa-Hattori <i>et al</i> , 1992)	
			Mancuso 2003 (Mancuso <i>et al</i> , 2003)	Parker 1989 (Parker <i>et al</i> , 1989)	

			Manczak 2004 (Manczak <i>et al.</i> , 2004)	Parker 2008 (Parker <i>et al.</i> , 2008)	
			Maurer 2000 (Maurer <i>et al.</i> , 2000)	Schaegger 1995 (Schägger, 1995)	
			Mutisya 1994 (Mutisya <i>et al.</i> , 1994)	Schapira 1990a (Schapira <i>et al.</i> , 1990a)	
			Parker 1990 (Parker <i>et al.</i> , 1990)	Schapira 1990b (Schapira <i>et al.</i> , 1990b)	
			Parker 1994a (Parker <i>et al.</i> , 1994b)	Shinde 2006 (Shinde and Pasupathy, 2006)	
			Parker 1994b (Parker <i>et al.</i> , 1994a)	Shoffner 1991 (Shoffner <i>et al.</i> , 1991)	
			Pérez-Gracia 2008 (Pérez-Gracia <i>et al.</i> , 2008)	Thomas 2012 (Thomas <i>et al.</i> , 2012)	
			Reichmann 1993 (Reichmann <i>et al.</i> , 1993)	Varghese 2009 (Varghese <i>et al.</i> , 2009)	
			Sekar 2015 (Sekar <i>et al.</i> , 2015)	Wiedemann 1999 (Wiedemann <i>et al.</i> , 1999)	
			Sheehan 1997 (Sheehan <i>et al.</i> , 1997)	Yoshino 1992 (Yoshino <i>et al.</i> , 1992)	
			Simonian 1993 (Simonian and Hyman, 1993)		
			Simonian 1995 (Simonian and Hyman, 1995)		
			Terni 2010 (Terni <i>et al.</i> , 2010)		
			Valla 2001 (Valla <i>et al.</i> , 2001)		
			Valla 2006 (Valla <i>et al.</i> , 2006)		
			Verwer 2000 (Verwer <i>et al.</i> , 2000)		
			Vitali 2009 (Vitali <i>et al.</i> , 2009)		
			Wong-Riley 1997 (Wong-Riley <i>et al.</i> , 1997)		


**Table 2. (Top) Heterogeneity and Inconsistency.** Significant Q statistics indicates the existence of heterogeneity. A  $I^2$  value of 0% indicates no observed inconsistency, whereas larger values shows increasing inconsistency. **(Bottom) Egger's regression test for funnel plot asymmetry.** The existence of funnel plot asymmetry and potential publication bias is indicated by p-values < 0.05. df = degrees of freedom.

<b>Heterogeneity and Inconsistency</b>				
		<b>Q</b>	<b>p-value</b>	<b><math>I^2</math> (%)</b>
<b>MDD</b>	<b>Complex I</b>	549.160	0.000	91%
	<b>Complex IV</b>	0.030	1.000	0%
<b>BD</b>	<b>Complex I</b>	137410.000	0.000	100%
	<b>Complex IV</b>	873.150	0.000	95%
<b>SZ</b>	<b>Complex I</b>	2197.500	0.000	97%
	<b>Complex IV</b>	209.560	0.000	88%
<b>AD</b>	<b>Complex I</b>	64812.000	0.000	100%
	<b>Complex IV</b>	4490.300	0.000	97%
<b>PD</b>	<b>Complex I</b>	437.650	0.000	80%
	<b>Complex IV</b>	3083.300	0.000	99%
<b>AGING</b>	<b>Complex I</b>	27.934	0.218	18%
	<b>Complex IV</b>	11.684	1.000	0%
<b>Egger's regression</b>				
		<b>z-value</b>	<b>df</b>	<b>p-value</b>
<b>MDD</b>	<b>Complex I</b>	-1.969	4	0.120
	<b>Complex IV</b>	0.630	2	0.593
<b>BD</b>	<b>Complex I</b>	-2.191	17	0.043
	<b>Complex IV</b>	-0.042	8	0.968
<b>SZ</b>	<b>Complex I</b>	0.135	13	0.895
	<b>Complex IV</b>	1.035	8	0.331
<b>AD</b>	<b>Complex I</b>	-0.109	17	0.915
	<b>Complex IV</b>	-1.381	37	0.176
<b>PD</b>	<b>Complex I</b>	-2.127	30	0.042
	<b>Complex IV</b>	-0.176	26	0.862
<b>AGING</b>	<b>Complex I</b>	-1.025	9	0.332
	<b>Complex IV</b>	-2.820	13	0.014









MDD	<ul style="list-style-type: none"> <li>• Small number of studies with small sample sizes</li> <li>• Region-specific heterogeneity</li> <li>• <b>Moderate effects in complex I, small effects in complex IV</b></li> </ul>
BPD	<ul style="list-style-type: none"> <li>• Small sample sizes</li> <li>• Region-specific heterogeneity</li> <li>• <b>Moderate effects in complex I, small effects in complex IV</b></li> </ul>
SZ	<ul style="list-style-type: none"> <li>• Small sample sizes</li> <li>• Region-specific heterogeneity</li> <li>• <b>Moderate effects in complex I and IV</b></li> </ul>
AD	<ul style="list-style-type: none"> <li>• Large number of studies with large sample sizes</li> <li>• Tissue-specific heterogeneity (low in blood, high in brain)</li> <li>• <b>Strong effects in complex I and IV</b></li> </ul>
PD	<ul style="list-style-type: none"> <li>• Large number of studies</li> <li>• No heterogeneity</li> <li>• <b>Strong effects in complex I and IV</b></li> </ul>
AGING	<ul style="list-style-type: none"> <li>• Most robust effects</li> <li>• No heterogeneity</li> <li>• <b>Strongest effects in complex I and IV</b></li> </ul>